

Product datasheet for **RC237206**

Thrombopoietin (THPO) (NM_001290027) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Thrombopoietin (THPO) (NM_001290027) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	THPO
Synonyms:	MGDF; MKCSF; ML; MPLLG; THCYT1; TPO
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	>RC237206 representing NM_001290027 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGGAGCTGACTGAATTGCTCCTCGTGGTCATGCTTCTCCTAACTGCAAGGCTAACGCTGTCCAGCCCGG
CTCCTCCTGCTTGTGACCTCCGAGTCCTCAGTAACTGCTTCGTGACTCCCATGTCCTTCACAGCAGACT
GAGCCAGTGCCAGAGGTTACCCCTTTGCCTACACCTGTCTGTGCCTGCTGTGGACTTTCAGTTGGGA
GAATGGAAAACCCAGATGGAGGAGACCAAGGCACAGGACATTCTGGGAGCAGTGACCCTTCTGCTGGAGG
GAGTGATGGCAGCACGGGGACAACCTGGGACCCACTTGCCTCTCATCCCTCCTGGGGCAGCTTTCTGGACA
GGTCCGTCCTCCTTGGGGCCCTGCAGAGCCTCCTTGGAAACCCAGCTTCTCCACAGGGCAGGACCACA
GCTCACAAGGATCCCAATGCCATCTTCTGAGCTTCCAACACCTGCTCCGAGGAAAGGACTTCTGGATTG
TTGGAGACAACTTCACTGCCTCAGCCAGAACTACTGGCTCTGGGCTTCTGAAGTGGCAGCAGGGATTCA
GAGCCAAGATTCTGGTCTGCTGAACCAAACCTCCAGGTCCTGGACCAATCCCCGGATACCTGAACAG
GATACACGAACCTTGAATGGAACCTGTGGACTCTTCTGGACCCTCACGCAGGACCCTAGGAGCCCCG
GACATTTCTCAGGAACATCAGACACAGGCTCCCTGCCACCAACCTCCAGCCTGGATATTCTCCTTCCC
CAACCCATCCTCCTACTGGACAGTATACGCTCTCCCTCTTCCACCCACCTTGCCACCCCTGTGGTCCA
GCTCCACCCCTGCTTCC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC237206 representing NM_001290027
 Red=Cloning site Green=Tags(s)

MELTELLLVMLLLTARLTLSSPAPPACDLRVLSKLLRDSHVLHSRLSQCEVHPLPTPVLLPAVDFSLG
 EWKTQMEETKAQDILGAVTLLLEGVMAARGQLGPTCLSSLLGQLSGQVRLLLGALQSLGTQLPPQGRIT
 AHKDPNAIFLSFQHLRKGDFWIVGDKLHCLSQNYWLWASEVAAGIQSQDSWSAEPNLQVPGPNRIPEQ
 DTRTLEWNSWTLSWTLTQDPRSPGHFLRNIRHRLPATQPPAWIFSFPNPSSYWTVYALPSSTHLAHPCGP
 APPPAS


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_001290027

ORF Size: 858 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001290027.1 , NP_001276956.1
RefSeq Size:	2074 bp
RefSeq ORF:	861 bp
Locus ID:	7066
UniProt ID:	P40225
Cytogenetics:	3q27.1
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Hematopoietic cell lineage
MW:	32 kDa
Gene Summary:	Megakaryocytopoiesis is the cellular development process that leads to platelet production. The main functional protein encoded by this gene is a humoral growth factor that is necessary for megakaryocyte proliferation and maturation, as well as for thrombopoiesis. This protein is the ligand for MLP/C_MPL, the product of myeloproliferative leukemia virus oncogene. Mutations in this gene are the cause of thrombocythemia 1. Alternative promoter usage and differential splicing result in multiple transcript variants differing in the 5' UTR and/or coding region. Multiple AUG codons upstream of the main open reading frame (ORF) have been identified, and these upstream AUGs inhibit translation of the main ORF at different extent. [provided by RefSeq, Feb 2014]