

## Product datasheet for RC237272

### SLAMF7 (NM\_001282592) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	SLAMF7 (NM_001282592) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	SLAMF7
Synonyms:	19A; CD319; CRACC; CS1
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	>RC237272 representing NM_001282592 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**GCGATCGCC**

ATGGCTGGTTCCCAACATGCCTCACCCCTCATCTATATCCTTTGGCAGCTCACAGGGTCAGCAGCCTCTG  
GACCCGTGAAAGAGCTGGTCGGTCCGTTGGTGGGCGGTGACTTCCCCCTGAAGTCAAAGTAAAGCA  
AGTTGACTCTATTGTCTGGACCTCAACACAACCCTCTTGTCACCATACAGCCAGAAGGGGGCACTATC  
ATAGTGACCCAAAATCGTAATAGGGAGAGAGTAGACTTCCCAGATGGAGGCTACTCCCTGAAGCTCAGCA  
AACTGAAGAAGAATGACTCAGGGATCTACTATGTGGGGATATACAGCTCATCACTCCAGCAGCCCTCCAC  
CCAGGAGTACGTGCTGCATGTCTACGAGCACCTGTCAAAGCCTAAAGTCACCATGGGTCTGCAGAGCAAT  
AAGAATGGCACCTGTGTGACCAATCTGACATGCTGCATGGAACATGGGGAAGAGGATGTGATTTATACCT  
GGAAGGCCCTGGGGCAAGCAGCCAATGAGTCCATAATGGGTCCATCCTCCCATCTCCTGGAGATGGGG  
AGAAAGTGATATGACCTTCATCTGCGTTGCCAGGAACCCTGTGAGCAGAACTTCTCAAGCCCCATCCTT  
GCCAGGAAGCTCTGTGAAGGTGCTGCTGATGACCCAGATTCCCATGGTCTCCTGTGTCTCCTGTTGG  
TGCCCTCCTGCTCAGTCTTTGTACTGGGGCTATTTCTTTGGTTTCTGAAGAGAGAGACAAGAAGA  
GAACAATCCTAAAGGAAGATCCAGCAAATACGGTTTACTCCACTGTGAAATACCGAAAAAGATGGAAAA  
TCCCCTCACTGCTCAGATGCCAGACACCAAGGCTATTTGCCTA

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

**Protein Sequence:** >RC237272 representing NM\_001282592  
 Red=Cloning site Green=Tags(s)

MAGSPTCLTIYILWQLTGSAAAGPVKELVGSVGGAVTFPLKSKVKQVDSIVWTFNTTPLVTIQPEGGTI  
 IVTQNRNRERVDFFDGGYSLKSLKKNDSGIYYVGIYSSSLQQPSTQEYVLHVYEHLSPKPKVTMGLQSN  
 KNGTCVTNLTCCEHGEEDVIYTKALGQAANESHNGSILPISWRWGESDMTFICVARNPVS RNFS SPIL  
 ARKLCEGAADDPDSSMVLCLLLVPLLLSLFVLGLFLWFLKRERQEENPKGRSSKYGLLHCGNTEKDGK  
 SPLTAHDARHTKAICL


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-MluI

**Cloning Scheme:**



**Plasmid Map:**



**ACCN:** NM\_001282592

**ORF Size:** 888 bp

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001282592.2</a>
<b>RefSeq Size:</b>	2804 bp
<b>RefSeq ORF:</b>	891 bp
<b>Locus ID:</b>	57823
<b>UniProt ID:</b>	<a href="#">Q9NQ25</a>
<b>Cytogenetics:</b>	1q23.3
<b>Protein Families:</b>	Druggable Genome, Transmembrane
<b>MW:</b>	33 kDa
<b>Gene Summary:</b>	Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2. Isoform 1 mediates NK cell activation through a SH2D1A-independent extracellular signal-regulated ERK-mediated pathway (PubMed:11698418). Positively regulates NK cell functions by a mechanism dependent on phosphorylated SH2D1B. Downstream signaling implicates PLCG1, PLCG2 and PI3K (PubMed:16339536). In addition to heterotypic NK cells-target cells interactions also homotypic interactions between NK cells may contribute to activation. However, in the absence of SH2D1B, inhibits NK cell function. Acts also inhibitory in T-cells (By similarity). May play a role in lymphocyte adhesion (PubMed:11802771). In LPS-activated monocytes negatively regulates production of proinflammatory cytokines (PubMed:23695528).[UniProtKB/Swiss-Prot Function]