

Product datasheet for RC237738

AZIN2 (NM_001301824) Human Tagged ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	AZIN2 (NM_001301824) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	AZIN2
Synonyms:	ADC; AZI2; AZIB1; ODC-p; ODC1L; ODCp
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>>RC237738 representing NM_001301824 Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C


ATGGAGTTGGTCCAGCATATTGGAATCCCTGCCAGTAAGATCATCTGCGCCAACCCCTGTAAGCAAATTG CACAGATCAAATATGCTGCCAAGCATGGGATCCAGCTGCTGAGCTTTGACAATGAGATGGAGCTGGCAAA GGTGGTAAAGAGCCACCCCAGTGCCAAGATGGTTCTGTGCATTGCTACCGATGACTCCCACTCCCTGAGC TGCCTGAGCCTAAAGTTTGGAGTGTCACTGAAATCCTGCAGACACCTGCTTGAAAATGCGAAGAAGCACC ATGTGGAGGTGGTGGGTGTGAGTTTTCACATTGGCAGTGGCTGTCCTGACCCTCAGGCCTATGCTCAGTC CATCGCAGACGCCCGGCTCGTGTTTGAAATGGGCACCGAGCTGGGTCACAAGATGCACGTTCTGGACCTT GGTGGTGGCTTCCCTGGCACAGAAGGGGCCCAAAGTGAGATTTGAAGAGATTGCTTCCGTGATCAACTCAG CCTTGGACCTGTACTTCCCAGAGGGCTGTGGCGTGGACATCTTTGCTGAGCTGGGGCGCTACTACGTGAC CTCGGCCTTCACTGTGGCAGTCAGCATCATTGCCAAGAAGGAGGTTCTGCTAGACCAGCCTGGCAGGGAG GAGGAAAATGGTTCCACCTCCAAGACCATCGTGTACCACCTTGATGAGGGCGTGTATGGGATCTTCAACT CAGTCCTGTTTGACAACATCTGCCCTACCCCCATCCTGCAGAAGAAACCATCCACGGAGCAGCCCCTGTA CAGCAGCAGCCTGTGGGGGCCCGGCGGTTGATGGCTGTGATTGCGTGGCTGAGGGCCTGTGGCTGCCGCAA CTACACGTAGGGGACTGGCTGGTCTTTGACAACATGGGCGCCTACACTGTGGGCATGGGTTCCCCCTTTT GGGGGACCCAGGCCTGCCACATCACCTATGCCATGTCCCGGGTGGCCTGGGAAGCGCTGCGAAGGCAGCT GATGGCTGCAGAACAGGAGGATGACGTGGAGGGTGTGTGCAAGCCTCTGTCCTGCGGCTGGGAGATCACA GACACCCTGTGCGTGGGCCCTGTCTTCACCCCAGCGAGCATCATG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG**GTTTAA**



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

QRIGENE AZIN2 (NM_001301824) Human Tagged ORF Clone - RC237738


Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 001301824.2</u>
RefSeq Size:	1849 bp
RefSeq ORF:	1098 bp
Locus ID:	113451
UniProt ID:	<u>Q96A70</u>
Cytogenetics:	1p35.1
Protein Families:	Druggable Genome
Protein Pathways:	Arginine and proline metabolism, Metabolic pathways
MW:	40.2 kDa

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Science AZIN2 (NM_001301824) Human Tagged ORF Clone – RC237738

The protein encoded by this gene belongs to the antizyme inhibitor family, which plays a role Gene Summary: in cell growth and proliferation by maintaining polyamine homeostasis within the cell. Antizyme inhibitors are homologs of ornithine decarboxylase (ODC, the key enzyme in polyamine biosynthesis) that have lost the ability to decarboxylase ornithine; however, retain the ability to bind to antizymes. Antizymes negatively regulate intracellular polyamine levels by binding to ODC and targeting it for degradation, as well as by inhibiting polyamine uptake. Antizyme inhibitors function as positive regulators of polyamine levels by sequestering antizymes and neutralizing their effect. This gene encodes antizyme inhibitor 2, the second member of this gene family. Like antizyme inhibitor 1, antizyme inhibitor 2 interacts with all 3 antizymes and stimulates ODC activity and polyamine uptake. However, unlike antizyme inhibitor 1, which is ubiquitously expressed and localized in the nucleus and cytoplasm, antizyme inhibitor 2 is predominantly expressed in the brain and testis and localized in the endoplasmic reticulum-golgi intermediate compartment. Recent studies indicate that antizyme inhibitor 2 is also expressed in specific cell types in ovaries, adrenal glands and pancreas, and in mast cells. The exact function of this gene is not known, however, available data suggest its role in cell growth, spermiogenesis, vesicular trafficking and secretion. Accumulation of antizyme inhibitor 2 has also been observed in brains of patients with Alzheimer's disease. There has been confusion in literature and databases over the nomenclature of this gene, stemming from an earlier report that a human cDNA clone (identical to ODCp/AZIN2) had arginine decarboxylase (ADC) activity (PMID:14738999). Subsequent studies in human and mouse showed that antizyme inhibitor 2 was devoid of arginine decarboxylase activity (PMID:19956990). Alternatively spliced transcript variants have been described for this gene. [provided by RefSeq, Sep 2014]

Product images:



Circular map for RC237738

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US