

Product datasheet for **RC237864**

Protein Kinase A regulatory subunit I alpha (PRKAR1A) (NM_001278433) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Protein Kinase A regulatory subunit I alpha (PRKAR1A) (NM_001278433) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Protein Kinase A regulatory subunit I alpha
Synonyms:	ACRDYS1; ADOHR; CAR; CNC; CNC1; PKR1; PPNAD1; PRKAR1; TSE1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RC237864 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGAGTCTGGCAGTACCGCCAGTGGAGGACGCAGCCTTCGAGAATGTGAGCTCTACGTCAGAG
 AGCATAACATTCAGCGCTGCTCAAAGATTCTATTGTGCAGTTGTGCACTGCTCGACCTGAGAGACCCAT
 GGATTCTCAGGGAATACTTTGAGAGGTTGGAGAAGGAGGAGGCAAAACAGATTCAGAATCTGCAGAAA
 GCAGGCACTCGTACAGACTCAAGGGAGGATGAGATTTCTCCTCCTCCACCCAACCCAGTGGTTAAAGGTA
 GGAGGCGACGAGGTGCTATCAGCGCTGAGGTCTACACGGAGGAAGATGCGGCATCCTATGTTAGAAAGGT
 TATACCAAAGATTACAAGACAATGGCCGCTTTAGCCAAAGCCATTGAAAAGAATGTGCTGTTTTACAT
 CTTGATGATAATGAGAGAAGTGATTTTTGATGCCATGTTTTCGGTCTCCTTTATCGCAGGAGAGACTG
 TGATTCAGCAAGGTGATGAAGGGGATAACTTCTATGTGATTGATCAAGGAGAGACGGATGTCTATGTTAA
 CAATGAATGGGCAACCAAGTGTGGGGAAGGAGGGAGCTTTGGAGAAGTCTTTGATTTATGGAACACCG
 AGAGCAGCCACTGTCAAAGCAAAGACAAATGTGAAATTGTGGGCATCGACCGAGACAGCTATAGAGAA
 TCCTCATGGGAAGCACACTGAGAAAGCGGAAGATGTATGAGGAATTCCTTAGTAAAGTCTCTATTTTAGA
 GTCTCTGGACAAGTGGGAACGTCTACGGTAGCTGATGCATTGGAACCAAGTGCAGTTTGAAGATGGGCAG
 AAGATTGTGGTGCAGGGAGAACCAGGGGATGAGTTCCTCATTATTTAGAGGGGTCAGCTGCTGTGCTAC
 AACGTCGGTGCAGAAAATGAAGAGTTTGTGAAGTGGGAAGATTGGGGCCTTCTGATTATTTTGGTGAAT
 TGCACTACTGATGAATCGTCTCGTGTGCCACAGTTGTGCTCGTGGCCCTTGAAGTGCCTTAAGCTG
 GACCGACCTAGATTTGAACGTGTTCTTGGCCATGCTCAGACATCCTCAAACGAAACATCCAGCAGTACA
 ACAGTTTTGTGCTACTGTCTGTC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>RC237864 protein sequence
 Red=Cloning site Green=Tags(s)

MESGSTAASEEARSLRECELYVQKHNIQALLKDSIVQLCTARPERPMAFLREYFERLEKEEAKQIQNLQK
 AGTRTDSREDEISPPPPNPVVKRRRRGAI SAEVYTEEDAASYVRKVI PKDYKTMALAKAIEKNVLF SH
 LDDNERSDIFDAMF SVSFIAGETVIQQGDEGDNFYVIDQGETDVYVNNWATS VEGGSGFELAL IYGT P
 RAATVKAKTNVKGIDRDSYRRILMGSTLRKRKMYEEFLSKVSILESLDKWERLTVADALEPVQFEDGQ
 KIVVQGPGEFFIILEGSAAVLQRRSENEEFVEVGR LGPSDYFGEIALLMNRPRAA TVVARGPLKCVKL
 DRPRFERVLGPCSDILKRNIQQYNSFVLSLV

TRTRPLEQKLI SEEDLAANDILDYKDDDDKV

Chromatograms:

https://cdn.origene.com/chromatograms/mk6013_d05.zip

Restriction Sites:

Sgfl-Mlul

Cloning Scheme:



ACCN: NM_001278433

ORF Size: 1143 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001278433.1](#), [NP_001265362.1](#)

RefSeq Size: 4328 bp

RefSeq ORF: 1146 bp

Locus ID: 5573

UniProt ID: [P10644](#)

Cytogenetics: 17q24.2


Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Apoptosis, Insulin signaling pathway

MW: 43 kDa

Gene Summary: cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-dependent protein kinase, which transduces the signal through phosphorylation of different target proteins. The inactive kinase holoenzyme is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. This gene encodes one of the regulatory subunits. This protein was found to be a tissue-specific extinguisher that down-regulates the expression of seven liver genes in hepatoma x fibroblast hybrids. Mutations in this gene cause Carney complex (CNC). This gene can fuse to the RET protooncogene by gene rearrangement and form the thyroid tumor-specific chimeric oncogene known as PTC2. A nonconventional nuclear localization sequence (NLS) has been found for this protein which suggests a role in DNA replication via the protein serving as a nuclear transport protein for the second subunit of the Replication Factor C (RFC40). Several alternatively spliced transcript variants encoding two different isoforms have been observed. [provided by RefSeq, Jan 2013]

Product images:



Circular map for RC237864