

Product datasheet for **RC238055**

PLAGL1 (NM_001289039) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PLAGL1 (NM_001289039) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	PLAGL1
Synonyms:	LOT1; ZAC; ZAC1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC238055 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCTACCCATTCTCCCAGAAATCTCACCAGTGTGCTCACTGTGAGAAGACGTTCAACCGAAAGACC
ACCTGAAAACCACCTCCAGACCCACGACCCCAACAAAATGGCCTTTGGGTGTGAGGAGTGTGGGAAGAA
GTACAACACCATGCTGGGCTATAAGAGGCACCTGGCCCTCCATGCGGCCAGCAGTGGGGACCTCACCTGT
GGGTCTGTGCCCTGGAGCTAGGGAGCACCGAGGTGCTACTGGACCACCTCAAAGCCCATCGGAAGAGA
AGCCCCCTAGCGAACCAAGGAAAAGAAGCACCAGTGCACCACCTGTGAAAGATGCTTCTACACCCGGAA
GGATGTGCGACGCCACCTGGTGGTCCACACAGGATGCAAGGACTTCTGTGCCAGTTCTGTGCCAGAGA
TTTGGGCGCAAGGATCACCTCACCCGGCATAACGAAGACCCACTCACAGGAGCTGATGAAAGAGAGCT
TGACAGACCGGAGACCTTCTGAGCACCTTCCACACCATCTCGCCTTCAATCCAACCTGAAGGCTGTGCCTT
GCCTCCTTTCCCTTAGGAGCTTCTGCCAGAACGGGCTTGAAGTAGCTTGCCAGCTGAGGTCCATAGC
CTCACCTCAGTCCCCAGAACAAGCCGCCAGCCTATGCAGCCGCTGCCAGAGTCCCTGGCCTCCCTCC
ACCCCTCGGTATCCCCTGGCTCTCTCCGCCACCCTTCCAATCACAAGTACAACACCACTTCTACCTC
ATACTCCCACCTGCAAGCCTGCCCTCAAAGCAGATACTAAAGTTTTTGCAATATCAGTTTGTGTTGAG
GACTTGCTCTGCAAGAGCCTCAGTCACCTCAAAGCTCAACCCAGGTTTTGATCTGGCTAAGGGAAATG
CTGGTAAAGTAAACCTGCCCAAGGAGCTGCCTGCAGATGCTGTGAACCTAACAATACCTGCCTCTGGA
CCTGTCCCCCTGTTGGGCTTCTGGCAGCTGCCCCCTCCTGTACCCAAAATACCTTTGGGAATAGCACT
CTTGCCCTGGGGCTGGGAATCTTTGCCCCACAGGTTAAGCTGTCTGGGGCAGCAGCAGCAAGAACCC
CACTTGCCATGGGCACTGTGAGCCTGGGCCAGCTCCCCCTGCCCCCATCCCTCATGTGTTCTCAGCTGG
CACTGGCTCTGCCATCCTGCCTCATTTCCATCATGCATTGAGA

ACGGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC238055 protein sequence
Red=Cloning site Green=Tags(s)

MATHSPQKSHQCAHCEKTFNRKDHLKNHLQTHDPNKMAFGCEEKGKYNMMLGYKRHLALHAASSGDLTC
 GVCALLEGSTEVLLDHLKAHAEEKPPSGTKEKKHQCDHCERCIFYTRKDVRRLVVHTGCKDFLCQFCAQR
 FGRKDHLTRHTKKTHSQELMKESLQTGDLLSTFHTISPSFQLKAAALPPFPLGASAQNGLASSLPAEVHS
 LTLSPPEQAAQPMQPLPESLASLHPSVSPGSPPPPLPNHKYNTTSTYSPLASLPLKADTKGFCNISLFE
 DLPLQEPQSPQKLNPGFDLAKGNAGKVNLPKELPADAVNLTIPASLDLSPLLGFWQLPPPATQNTFGNST
 LALGPGESLPHRLSCLGQQQEPPPLAMGTVSLGQLPLPPIPHVF SAGTGSAILPHFHHAFR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6446_h09.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001289039

ORF Size: 1233 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001289039.2](#)

RefSeq Size: 2760 bp

RefSeq ORF: 1236 bp

Locus ID: 5325

UniProt ID: [Q9UM63](#)


Cytogenetics: 6q24.2

Protein Families: Transcription Factors

MW: 44.7 kDa

Gene Summary: This gene encodes a C2H2 zinc finger protein that functions as a suppressor of cell growth. This gene is often deleted or methylated and silenced in cancer cells. In addition, overexpression of this gene during fetal development is thought to be the causal factor for transient neonatal diabetes mellitus (TNDM). Alternative splicing and the use of alternative promoters results in multiple transcript variants encoding two different protein isoforms. The P1 downstream promoter of this gene is imprinted, with preferential expression from the paternal allele in many tissues. [provided by RefSeq, Nov 2015]

Product images:



Circular map for RC238055