

Product datasheet for **RC239757**

eIF3s8 (EIF3C) (NM_001286478) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	eIF3s8 (EIF3C) (NM_001286478) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	eIF3s8
Synonyms:	eIF3-p110; EIF3CL; EIF3S8
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide
Sequence:

>RC239757 representing NM_001286478
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGCATCGC**

ATGTCGCGGTTTTTCACCAACCGTTTCGGACAGCGAGTCCGAGTCGTCCTTGTCGGGGAGGAGCTCGTCA
CCAAACCTGTGCGAGGCAACTATGGCAAACAGCCATTGTTGCTGAGCGAGGATGAAGAAGATACCAAGAG
AGTTGTCCGAGTGCCAAGGACAAGAGTTTGAGGAGCTGACCAACCTTATCCGGACCATCCGTAATGCC
ATGAAGATTCTGATGTACCAAGTGCCTGGAAGAGTTTGAGCTCCTGGGAAAAGCATATGGGAAGGCCA
AAAGCATTGTGGACAAAGAAGGTGTCCCCGGTTCTATATCCGCATCCTGGCTGACCTAGAGGACTATCT
TAATGAGATGAACAAGAACAATGCCAAGGCTCTGAGCACCTTGCGTCAGAAGATCCGAAAATACAACCGT
GATTTGAGTCCCATATCACAAGCTACAAGCAGAACCCGAGCAGTCTGCGGATGAAGATGCTGAGAAAA
ATGAGGAGGATTGAGAAGGCTCTCAGATGAGGATGAGGATGAGGACGGAGTCAGTGCTGCAACTTTCTT
GAAGAAGAAATCAGAAGCTCCTTCTGGGAGAGTCGCAAGTTCTCAAAAAGATGGATGATGAAGATGAG
GACTCAGAAGATTCCGAAGATGATGAAGACTGGGACACAGGTTCCACATCTTCCGATTCCGACTCAGAGG
AGGAAGAAGGGAAACAAACCGCGCTGGCCTCAAGATTTCTTAAAAAGGCACCCACCAAGATGAGGACAA
GAAGGCAGCCGAGAAGAAACGGGAGGACAAAGCTAAGAAGAAGCACGACAGGAAATCCAAGCGCCTGGAT
GAGGAGGAGGAGGACAATGAAGGCGGGAGTGGGAAAGGTCGCGGGCGGAGTGCCGTGGTTAAGGAGA
AGCCAAAATGTTTGCCAAAGGGAAGTGAATCACCATGCTGTTGTTATCAAGAACTGAATGAGATCCT
ACAGGCACGAGGCAAGAAGGGAAGTATGCTGCTGCCAGATTGAGCTGCTGCAACTGCTGGTTGAGATT
GCAGCGGAAAACAACCTGGGAGAGGGCGTCAATGTCAAGATCAAGTTCAATATCATCGCCTCTCTCTATG
ACTACAACCCCAACCTGGCAACCTACATGAAGCCAGAGATGTGGGGAAGTGCCTGGACTGCATCAATGA
GCTGATGGATATCCTGTTTGCAAAATCCCAACATTTTGTGGAGAGAATATTCTGGAAGAGAGTGGAAC
CTGCACAACGCTGACCAGCCACTGCGTGTCCGTGGCTGCATCCTAACTCTGGTGGAAAGATGGATGAAG
AATTTACCAAAATATGCAAAATACTGACCCTCACTCCCAAGAGTACGTGGAGCACTTGAAGGATGAGGC
CCAGGTGTGTCCATCATCGAGCGTGTGACGCGTACCTGGAGGAGAAGGGCACTACCGAGGAGGTCTGC
CGCATCTACCTGCTGCGCATCCTGCACACCTACTACAAGTTTGATTACAAGGCCCATCAGCGACAGCTGA
CCCCGCTGAGGGCTCCTCAAAGTCTGAGCAAGACCAGGCAGAAAATGAGGGCGAGGACTCGGCTGTGTT
GATGGAGAGACTGTGCAAGTACATCTACGCCAAGGACCGCACAGACCGGATCCGCACATGTGCCATCCTC
TGCCACATCTACCACCATGCTCTGCACTCGCGCTGGTACCAGGCCGCGACCTCATGCTCATGAGCCACT
TGCAAGACAACATTCAGCATGCAGACCCGCCAGTGCAGATCCTTTACAACCGCACCATGGTGCAGCTGGG
CATCTGTGCCTTCCGCCAAGGCCTGACCAAGGACGCACACAACGCCCTGCTGGACATCCAGTCGAGTGGC
CGAGCCAAGGAGCTTCTGGGCCAGGGCCTGCTGCTGCGCAGCCTGCAGGAGCGCAACCAGGAGCAGGAGA
AGGTGGAGCGGCGCCGTGAGTCCCCCTCCACCTGCACATCAACCTGGAGCTGCTGGAGTGTGCTACCT
GGTGTCTGCCATGCTCCTGGAGATCCCCTACATGGCCGCCCATGAGAGCGATGCCCGCCGACGCATGATC
AGCAAGCAGTTCACCAACAGCTGCGCGTGGGCGAGCGACAGCCCTGCTGGGTCCCCCTGAGTCCATGC
GGGAACATGTGGTCGCTGCCTCCAAGGCCATGAAGATGGGTGACTGGAAGACCTGTCACAGTTTATCAT
CAATGAGAAGATGAATGGGAAAGTGTGGGACCTTTTCCCGAGGCTGACAAAGTCCGCACCATGCTGGTT
AGGAAGATCCAGGAAGAGTCACTGAGGACCTACCTCTTACCTACAGCAGTGTCTATGACTCCATCAGCA
TGGAGACGCTGTGAGACATGTTTGAGCTGGATCTGCCCACTGTGCACTCCATCATCAGCAAAATGATCAT
TAATGAGGAGCTGATGGCCTCCCTGGACCAGCCAACACAGACAGTGGTATGCACCGCACTGAGCCCACT
GCCAGCAGAACCTGGCTCTGCAGCTGGCCGAGAAGCTGGGCAGCCTGGTGGAGAACAACGAACGGGTGT
TTGACCACAAGCAGGGCACCTACGGGGCTACTTCCGAGACCAGAAGGACGGCTACCGCAAAACGAGGG
CTACATGCGCCGCGGTGGCTACCGCCAGCAGCAGTCTCAGACGGCCTAC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC239757 representing NM_001286478
Red=Cloning site Green=Tags(s)


MSRFFTTGSDSESSLSGEELVTKPVGGNYGKQPLLLSEDEEDTKRVVRSKDKRFEELTNLIRTIRNA
MKIRDVTKCLEEFELLGKAYGKAKSIVDKEGVPRFYIRILADLEDYLNEMNKNNAKALSTLRQKIRKYNR
DFESHITSYKQNPESADEDAEKNEEDSEGSSDEDEDEDGVSAAFLKKKSEAPSGESRKFLKKMDEDE
DSEDEDEDEDWDTGSTSSSDSDEEEEGQTALASRFLKKAPTDDKAAEKKREDKAKKKHDKSKRLD
EEEEEDNEGGEWERVGGVPLVKEKPKMFAKGTEITHAVVIKKLNEILQARGKKGTDRAAQIELLQLLVQI
AAENNLGEGVIVKIKFNIIASLYDYNPNLATYMKPEMWGKCLDCINELMDILFANPNIFVGENILEESEN
LHNADQPLRVGRCILTLVERMDEEFTKIMQNTDPHSQEYVEHLKDEAQVCAIIERVQRYLEEKGTTEEVC
RIYLLRILHTYYKFDYKAHQRLTPPEGSSKSEQDAENEGEDSAVLMERLCKYIYAKDRTDRIRTCAIL
CHYHHALHSRWYQARDLMLMSHLQDNIQHADPPVQILYNRTMVQLGICAFRQGLTKDAHNALLDIQSSG
RAKELLGQGLLLRSLQERNQEQEKVERRRQVPFHLHINLELLECVYLVSAMLLEIPYMAAHESDARRMI
SKQFHHLRVGERQPLLGPPEMREHVVAASKAMKMGDWKTCHSFINEKMNGKVWDLFPEADKVRTMLV
RKIQEESLRTYLFYSSVYDSISMETLSDMFELDLPTVHSIISKMIINEELMASLDQPTQTVMHRTEPT
AQQNLALQLAEKLGSLVENNERVFDHKQGTYYGGYFRDQKDGYNKGYMRRGGYRQQSQATAY

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:




ACCN: NM_001286478

ORF Size:	2709 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_001286478.1 , NP_001273407.1
RefSeq Size:	3066 bp
RefSeq ORF:	2712 bp
Locus ID:	8663
UniProt ID:	Q99613
Cytogenetics:	16p11.2
Protein Families:	Druggable Genome
MW:	104.6 kDa

Gene Summary:

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:17581632, PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA_i and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773).[UniProtKB/Swiss-Prot Function]

Product images:


Circular map for RC239757