

Product datasheet for **RC400100**

Isocitrate dehydrogenase (IDH1) (NM_005896) Human Mutant ORF Clone

Product data:

Product Type:	Mutant ORF Clones
Product Name:	Isocitrate dehydrogenase (IDH1) (NM_005896) Human Mutant ORF Clone
Mutation Description:	R132G
Affected Codon#:	132
Affected NT#:	c. 394
Nucleotide Mutation:	IDH1 mutant (R132G), Myc-DDK-tagged ORF clone of Homo sapiens isocitrate dehydrogenase 1 (NADP+), soluble (IDH1) as transfection-ready DNA
Effect:	Missense
Symbol:	Isocitrate dehydrogenase
Synonyms:	HEL-216; HEL-S-26; IDCD; IDH; IDP; IDPC; PICD
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_005896
ORF Size:	1245 bp
Restriction Sites:	Sgfl-Mlul



[View online »](#)

ORF Nucleotide Sequence:

>RC400100 representing NM_005896
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**GCGATCGCC**

ATGTCCAAAAAATCAGTGGCGTTCTGTGGTAGAGATGCAAGGAGATGAAATGACACGAATCATTGGG
 AATTGATTAAAGAGAACTCATTTCCTACGTGGAATTGGATCTACATAGCTATGATTAGGCATAGA
 GAATCGTGATGCCACCAACGACCAAGTACCAAGGATGCTGCAGAAGCTATAAAGAAGCATAATGTTGGC
 GTCAAATGTGCCACTATCACTCCTGATGAGAAGAGGGTTGAGGAGTTCAAGTTGAAACAAATGTGGAAT
 CACCAATGGCACCATACGAAATATTCTGGGTGGCAGGCTCTCAGAGAAGCCATTATCTGCAAAAATAT
 CCCCCGGCTTGAGTGGATGGTAAAACCTATCATCATAGGTGGTCATGCTTATGGGGATCAATACAGA
 GCAACTGATTTTGTGTTCTCTGGCCTGAAAAGTAGAGATAACCTACACACCAAGTACGGAACCCAAA
 AGGTGACATACCTGGTACATAACTTTGAAGAAGTGGTGGTGTGCCATGGGGATGTATAATCAAGATAA
 GTCAATTGAAGATTTTGCACACAGTTCCTTCCAATGGCTCTGTCTAAGGGTTGGCCTTTGTATCTGAGC
 ACCAAAAACACTATTCTGAAGAAATATGATGGGCGTTTTAAAGACATCTTTCAGGAGATATATGACAAGC
 AGTACAAGTCCCAGTTTGAAGCTCAAAAGATCTGGTATGAGCATAGGCTCATCGACGACATGGTGGCCCA
 AGCTATGAAATCAGAGGGAGGCTTCATCTGGGCTGTAAAAACTATGATGGTGACGTGCAGTCCGACTCT
 GTGGCCCAAGGGTATGGCTCTCTCGGCATGATGACCAGCGTGTGGTTTGTCCAGATGGCAAGACAGTAG
 AAGCAGAGGCTGCCACGGGACTGTAACCCGCTCACTACCGCATGTACCAGAAAGGACAGGAGACGTCCAC
 CAATCCCATTGCTTCCATTTTGCCTGGACCAGAGGGTTAGCCACAGAGCAAAGCTTGATAACAATAAA
 GAGCTTGCCTTCTTTCAAATGCTTTGGAAGAAGTCTCTATTGAGACAATTGAGGCTGGCTTCATGACCA
 AGGACTTGGCTGCTTGCAATAAAGGTTTACCCAATGTGCAACGTTCTGACTACTTGAATACATTTGAGTT
 CATGGATAAACTTGGAGAAAACCTGAAGATCAAACCTAGCTCAGGCCAAACTT

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>RC400100 representing NM_005896
 Red=Cloning site Green=Tags(s)


MSKKISGGSVEMQGDDEMTRIIWELIKEKLIFPYVELDLHSYDLGIENRDATNDQVTKDAAEAIKKNHVG
 VKCATITPDEKRVEEFKLMWQKSPNGTIRNILGGTVFREAIICKNIPRLVSGWVKPIIIGGHAYGDQYR
 ATDFVVPGPVKVEITYTPSDGTQKVTYLVHNFEEGGVAMGMYNQDKSIEDFAHSSFQMALSKGWPLYLS
 TKNTILKKYDGRFKDIFQEIYDKQYKSQFEAQKIWYEHRLIDDMVAQAMKSEGGFIWACKNYDGDVQSDS
 VAQGYGSLGMMTSVLVCPDGKTVEAAHGTVTRHYRMYQKGQETSTNPIASIFAWTRGLAHRAKLDNNK
 ELAFFANALEEVSIEIIEAGFMTKDLAACIKGLPNVQRSDYLNTEFMDKLGENLKIKLAQAKL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-Mlul

Cloning Scheme:



OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in *E. coli* are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

RefSeq:

[NP_005887](#)

RefSeq Size:

2339 bp

RefSeq ORF:

1245 bp

Locus ID:

3417

Cytogenetics:

2q34

Domains:

isodh

Protein Pathways:

Citrate cycle (TCA cycle), Glutathione metabolism, Metabolic pathways

MW:

46.5 kDa

Gene Summary:

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Sep 2013]