

Product datasheet for **RC400449**

PTEN (NM_000314) Human Mutant ORF Clone

Product data:

Product Type:	Mutant ORF Clones
Product Name:	PTEN (NM_000314) Human Mutant ORF Clone
Mutation Description:	E235*
Affected Codon#:	235
Affected NT#:	c.703
Nucleotide Mutation:	PTEN Mutant (E235*), Myc-DDK-tagged ORF clone of Homo sapiens phosphatase and tensin homolog (PTEN) as transfection-ready DNA
Effect:	Truncation
Symbol:	PTEN
Synonyms:	10q23del; BZS; CWS1; DEC; GLM2; MHAM; MMAC1; PTEN1; PTENbeta; TEP1
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_000314
ORF Size:	702 bp
Restriction Sites:	Sgfl-Mlul



[View online »](#)

ORF Nucleotide Sequence:

>RC400449 representing NM_000314
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**GCGATCGCC**

ATGACAGCCATCATCAAAGAGATCGTTAGCAGAAACAAAAGGAGATATCAAGAGGATGGATTCGACTTAG
 ACTTGACCTATATTTATCCAAACATTATTGCTATGGGATTTCTGCAGAAAGACTTGAAGCGTATACAG
 GAACAATATTGATGATGTAGTAAGGTTTTGGATTCAAAGCATAAAAACCATTACAAGATATACAATCTT
 TGTGCTGAAAGACATTATGACACCGCCAAATTTAATTGCAGAGTTGCACAATATCCTTTTGAAGACCATA
 ACCCACCACAGCTAGAACTTATCAAACCCTTTGTGAAGATCTTGACCAATGGCTAAGTGAAGATGACAA
 TCATGTTGCAGCAATTCAGTGTAAAGCTGAAAAGGGACGAACTGGTGAATGATATGTGCATATTTATTA
 CATCGGGGCAATTTTTAAAGGCACAAGAGGCCCTAGATTTCTATGGGAAGTAAAGACCAGAGACAAAA
 AGGGAGTAACTATTCCAGTCAGAGGCGCTATGTGATTATTATAGCTACCTGTTAAAGAATCATCTGGA
 TTATAGACCAGTGGCACTGTTGTTTACAAGATGATGTTTGAAGTATTCCAATGTTCAAGTGGCGGAAGT
 TGCAATCCTCAGTTTGTGGTCTGCCAGCTAAAGGTGAAGATATATTCCTCCAATTCAGGACCCACACGAC
 GG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>RC400449 representing NM_000314
 Red=Cloning site Green=Tags(s)


MTAIIKEIVSRNKRRYQEDGFDDLTYIYPNIIAMGFPAERLEGVYRNNIDDVVRFLDSKHKNHYKIYNL
 CAERHYDTAKFNCRVAQYPFEDHNPPQLELIKPFCELDQWLSDDNHVAAIHCKAGKGRGVMICAYLL
 HRGKFLKAQEALDFYGEVTRDKKGVIPSQRVYVYYSYLLKNHLDYRVPVALLFHKMMFETIPMFSGGT
 CNPQFVVCQLKVKIYSSNSGPTRR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
RefSeq:	NP_000305
RefSeq Size:	5572 bp
RefSeq ORF:	1212 bp
Locus ID:	5728
Cytogenetics:	10q23.31
Domains:	PTPc_motif
Protein Families:	Druggable Genome, Phosphatase
Protein Pathways:	Endometrial cancer, Focal adhesion, Glioma, Inositol phosphate metabolism, Melanoma, p53 signaling pathway, Pathways in cancer, Phosphatidylinositol signaling system, Prostate cancer, Small cell lung cancer, Tight junction
MW:	27 kDa

Gene Summary:

This gene was identified as a tumor suppressor that is mutated in a large number of cancers at high frequency. The protein encoded by this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway. The use of a non-canonical (CUG) upstream initiation site produces a longer isoform that initiates translation with a leucine, and is thought to be preferentially associated with the mitochondrial inner membrane. This longer isoform may help regulate energy metabolism in the mitochondria. A pseudogene of this gene is found on chromosome 9. Alternative splicing and the use of multiple translation start codons results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2015]