

Product datasheet for **RC401282**

Glucokinase (GCK) (NM_000162) Human Mutant ORF Clone

Product data:

Product Type:	Mutant ORF Clones
Product Name:	Glucokinase (GCK) (NM_000162) Human Mutant ORF Clone
Mutation Description:	E248X
Affected Codon#:	248
Affected NT#:	742
Nucleotide Mutation:	GCK Mutant (E248X), Myc-DDK-tagged ORF clone of Homo sapiens glucokinase (hexokinase 4) (GCK), transcript variant 1 as transfection-ready DNA
Effect:	Diabetes, MODY
Symbol:	GCK
Synonyms:	FGQTL3; GK; GLK; HHF3; HK4; HKIV; HXKP; LGLK; MODY2; PNDM1
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_000162
ORF Size:	741 bp
Restriction Sites:	Sgfl-MluI



[View online »](#)

ORF Nucleotide Sequence:

>RC401282 representing NM_000162
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGGATCGCC

ATGCTGGACGACAGCCAGGATGGAGGCCCAAGAAGGAGAAGGTAGAGCAGATCCTGGCAGAGTTCC
 AGCTGCAGGAGGAGACCTGAAGAAGGTGATGAGACGGATGCAGAAGGAGATGGACCGCGCCTGAGGCT
 GGAGACCCATGAAGAGGCCAGTGTGAAGATGCTGCCACCTACGTGCGCTCCACCCAGAAAGGCTCAGAA
 GTCGGGGACTTCTCCTGGACCTGGGTGGCACTAACTTCAGGGTGTGCTGGTGAAGGTGGGAGAAG
 GTGAGGAGGGCAGTGGAGCGTGAAGACCAAACACCAGATGACTCCATCCCGAGGACGCCATGACCGG
 CACTGCTGAGATGCTCTCGACTACATCTCTGAGTGCATCTCCGACTTCTGGACAAGCATCAGATGAAA
 CACAAGAAGCTGCCCTGGGCTTCACTTCTCCTTTCCTGTGAGGCACGAAGACATCGATAAGGGCATCC
 TTCTCAACTGGACCAAGGGCTTCAAGGCCTCAGGAGCAGAAGGGAACAATGTCGTGGGGCTTCTGCGAGA
 CGCTATCAAACGGAGAGGGGACTTTGAAATGGATGTGGTGGCAATGGTGAATGACACGGTGGCCACGATG
 ATCTCCTGCTACTACGAAGACCATCAGTGCAGGTCGGCATGATCGTGGGCACGGCTGCAATGCCTGCT
 ACATGGAGGAGATGCAGAATGTGGAGCTGGTGGAGGGGAC

AGCGGACCGACGCGTACGCGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC
 TGGATTACAAGGATGACGACGA TAAGGTTTAA

Protein Sequence:

>RC401282 representing NM_000162
 Red=Cloning site Green=Tags(s)


MLDDRARMEAAKKEKVEQILAEFQLQEEDLKKVMRRMQKEMDRGLRLETHEEASVKMLPTYVVRSTPEGSE
 VGDFLSLDLGGTNFRVMLVKVGEEGQWSVKTKHQMSIPEDAMTGAEMLFDYISECISDFLDKHQMK
 HKKLPLGFTTFFPVRHEDIDKILLNWTGFKASGAEGNNAVGLLRDAIKRRGDFEMDVVAMVNDTVATM
 ISCYYEDHQCEVGMIVGTGCNACymeemQNVELVEGD

SGPTRRRRLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
RefSeq:	NP_000153
RefSeq Size:	741 bp
RefSeq ORF:	1398 bp
Locus ID:	2645
Cytogenetics:	7p13
Protein Families:	Druggable Genome
Protein Pathways:	Amino sugar and nucleotide sugar metabolism, Galactose metabolism, Glycolysis / Gluconeogenesis, Insulin signaling pathway, Maturity onset diabetes of the young, Metabolic pathways, Starch and sucrose metabolism, Type II diabetes mellitus
MW:	27.2 kDa
Gene Summary:	This gene encodes a member of the hexokinase family of proteins. Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. In contrast to other forms of hexokinase, this enzyme is not inhibited by its product glucose-6-phosphate but remains active while glucose is abundant. The use of multiple promoters and alternative splicing of this gene result in distinct protein isoforms that exhibit tissue-specific expression in the pancreas and liver. In the pancreas, this enzyme plays a role in glucose-stimulated insulin secretion, while in the liver, this enzyme is important in glucose uptake and conversion to glycogen. Mutations in this gene that alter enzyme activity have been associated with multiple types of diabetes and hyperinsulinemic hypoglycemia. [provided by RefSeq, Aug 2017]