

Product datasheet for RC402270

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

G protein alpha S (GNAS) (NM_000516) Human Mutant ORF Clone

Product data:

Product Type: Mutant ORF Clones

Product Name: G protein alpha S (GNAS) (NM_000516) Human Mutant ORF Clone

Mutation Description: W154X

Affected Codon#: 154

Affected NT#: 462

Nucleotide Mutation: GNAS Mutant (W154X), Myc-DDK-tagged ORF clone of Homo sapiens GNAS complex locus

(GNAS), transcript variant 1 as transfection-ready DNA

Effect: Pseudohypoprhyroidism 1

Symbol: GNAS

Synonyms: AHO; C20orf45; GNAS1; GPSA; GSA; GSP; NESP; PITA3; POH; SCG6; SgVI

E. coli Selection: Kanamycin (25 ug/mL)

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001)

Tag: Myc-DDK
ACCN: NM 000516

ORF Size: 459 bp

Restriction Sites: Sgfl-Notl

G protein alpha S (GNAS) (NM_000516) Human Mutant ORF Clone - RC402270

ORF Nucleotide Sequence:

>RC402270 representing NM_000516

Red=Cloning site Blue=ORF Green=Tags(s)

ATGGGCTGCCTCGGGAACAGTAAGACCGAGGACCAGCGCAACGAGGAGAAGGCGCAGCGTGAGGCCAACA
AAAAGATCGAGAAGCAGCTGCAGAAGGACAAGCAGGTCTACCGGGCCACCGCCTGCTGCTGCTGGG
TGCTGGAGAATCTGGTAAAAGCACCATTGTGAAGCAGATGAGGATCCTGCATGTTAATGGGTTTAATGGA
GAGGGCGGCGAAGAGGACCCGCAGGCTGCAAGGAGCAACAGCGATGGTGAGAAGGCAACCAAAGTGCAGG
ACATCAAAAACAACCTGAAAGAGGCGATTGAAACCATTGTGGCCGCCATGAGCAACCTGGTGCCCCCCGT
GGAGCTGGCCAACCCCGAGAACCAGTTCAGAGTGGACTACATCCTGAGTGTGATGAACGTGCCTGACTTT
GACTTCCCTCCCGAATTCTATGAGCATGCCAAGGCTCTG

AGCGGACCGACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTCGGATTACAAGGATGACGACGA TAAGGTTTAA

Protein Sequence:

>RC402270 representing NM_000516 Red=Cloning site Green=Tags(s)

MGCLGNSKTEDQRNEEKAQREANKKIEKQLQKDKQVYRATHRLLLLGAGESGKSTIVKQMRILHVNGFNG EGGEEDPQAARSNSDGEKATKVQDIKNNLKEAIETIVAAMSNLVPPVELANPENQFRVDYILSVMNVPDF DFPPEFYEHAKAL

SGPTRTRRLEQKLISEEDLAANDILDYKDDDDK**V**


Restriction Sites:

Cloning Scheme:

Cloning sites used for ORF Shuttling:

Sgfl-Notl





^{*} The last codon before the Stop codon of the ORF



OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

RefSeq: <u>NP 000507</u>

 RefSeq Size:
 459 bp

 RefSeq ORF:
 1185 bp

 Locus ID:
 2778

Cytogenetics: 20q13.32

Domains: G-alpha


Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Calcium signaling pathway, Dilated cardiomyopathy, Gap junction, GnRH signaling pathway,

Long-term depression, Melanogenesis, Taste transduction, Vascular smooth muscle

contraction, Vibrio cholerae infection

MW: 16.8 kDa





Gene Summary:

This locus has a highly complex imprinted expression pattern. It gives rise to maternally, paternally, and biallelically expressed transcripts that are derived from four alternative promoters and 5' exons. Some transcripts contain a differentially methylated region (DMR) at their 5' exons, and this DMR is commonly found in imprinted genes and correlates with transcript expression. An antisense transcript is produced from an overlapping locus on the opposite strand. One of the transcripts produced from this locus, and the antisense transcript, are paternally expressed noncoding RNAs, and may regulate imprinting in this region. In addition, one of the transcripts contains a second overlapping ORF, which encodes a structurally unrelated protein - Alex. Alternative splicing of downstream exons is also observed, which results in different forms of the stimulatory G-protein alpha subunit, a key element of the classical signal transduction pathway linking receptor-ligand interactions with the activation of adenylyl cyclase and a variety of cellular reponses. Multiple transcript variants encoding different isoforms have been found for this gene. Mutations in this gene result in pseudohypoparathyroidism type 1a, pseudohypoparathyroidism type 1b, Albright hereditary osteodystrophy, pseudopseudohypoparathyroidism, McCune-Albright syndrome, progressive osseus heteroplasia, polyostotic fibrous dysplasia of bone, and some pituitary tumors. [provided by RefSeq, Aug 2012]