

Product datasheet for **RC402344**

Tuberin (TSC2) (NM_000548) Human Mutant ORF Clone

Product data:

Product Type:	Mutant ORF Clones
Product Name:	Tuberin (TSC2) (NM_000548) Human Mutant ORF Clone
Mutation Description:	Q373X
Affected Codon#:	373
Affected NT#:	1117
Nucleotide Mutation:	TSC2 Mutant (Q373X), Myc-DDK-tagged ORF clone of Homo sapiens tuberous sclerosis 2 (TSC2), transcript variant 1 as transfection-ready DNA
Effect:	Tuberous sclerosis
Symbol:	Tuberin
Synonyms:	LAM; PPP1R160; TSC4
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_000548
ORF Size:	1116 bp
Restriction Sites:	SgfI-XhoI



[View online »](#)

ORF Nucleotide Sequence:

>RC402344 representing NM_000548
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGCCAAACCAACAAGCAAAGATTAGGCTTGAAGGAGAAGTTAAGATTCTGTGGGACTGGGAACAC
 CGAGGCCAAATCCCAGGTCTGCAGAGGGTAAACAGACGGAGTTATCATCACCGCGAAATACTGAGAGA
 ACTGAGCATGGAATGTGGCCTCAACAATCGCATCCGGATGATAGGGCAGATTTGTGAAGTCGAAAAACC
 AAGAAATTTGAAGAGCACGCAGTGAAGCACTCTGGAAGGCGGTGCGGGATCTGTTGCAGCCGGAGCGGC
 CGCTGGAGGCCCGGCACGCGGTCTGGCTCTGCTGAAGGCCATCGTGCAGGGCAGGGCAGCGTTTGGG
 GGTCTCAGAGCCCTCTCTTTAAGGTCATCAAGGATTACCTTCCAACGAAGACCTTACGAAAGGCTG
 GAGTTTTCAAGGCCCTCACAGACAATGGGAGACACATCACCTACTTGGAGGAAGAGCTGGCTGACTTTG
 TCCTGCAGTGGATGGATGTTGGCTTGTCTCGGAATTCCTTCTGGTGTGGTGAACCTGGTCAAATCAA
 TAGCTGTTACCTCGACGAGTACATCGCAAGGATGGTTCAGATGATCTGTCTGCTGTGCGTCCGGACCGCG
 TCCTCTGTGGACATAGAGGTCTCCCTGCAGGTCTGGACGCCGTGGTCTGCTACAACCTGCCTGCCGGCTG
 AGAGCCTCCCGCTGTTATCGTTACCCTCTGTGCAACCATCAACGTCAAGGAGCTCTGCGAGCCTTGCTG
 GAAGCTGATGCGGAACCTCCTTGGCACCCACCTGGGCCACAGCGCCATCTACAACATGTGCCACCTCATG
 GAGGACAGAGCCTACATGGAGGACGCGCCCTGCTGAGAGGAGCCGTGTTTTTGTGGGCATGGCTCTCT
 GGGGAGCCACCGGCTCTATTCTCTCAGGAACCTGCCGACATCTGTGTTGCCATCATTTTACCAGGCCAT
 GGCATGTCCGAACGAGGTGGTGTCTATGAGATCGTCTGTCCATCACCAGGCTCATCAAGAAGTATAGG
 AAGGAGCTCCAGGTGGTGGCGTGGGACATTCTGCTGAACATCATCGAACGGCTCCTTACGACGCTC

AG**CGGACCG**ACGCGTACGCGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC
 TGGATTACAAGGATGACGACGA TAAGGTTTAA

Protein Sequence:


>RC402344 representing NM_000548
 Red=Cloning site Green=Tags(s)

MAKPTSKDSGLKEFKILLGLGTPRPNRPSAEGKQTEFIITAEILRELSMECGLNRRIRMIQICEVAKT
 KKFEHAEALWKAVADLLQPERPLEARHAVLALLKAIVQGGERLGVLRALFFKVIKDYPSNEDLHERL
 EVFKALTDNGRHITYLEEELADFVLQWMDVGLSSEFLLVLNLFKFNVCYLDEYIARMVQMICLLCVRTA
 SSVDIEVSLQVLDVAVCYNCLPAESLPLFIVTLCRTINVKELCEPCWKLNRNLLGTHLGHSAIYNMCHLM
 EDRAVMEDAPLLRGAVFFVGMALWGAHRLYSLRNSPTSVLPSFYQAMACPNEVVSYEIVLSITRLIKKYR
 KELQVVAWDILLNIIERLLQQL

SGP**TRRRLEQKLI**SEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-XhoI

Cloning Scheme:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Note:

Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq:

[NP_000539](#)

RefSeq Size:

1116 bp

RefSeq ORF:

5424 bp

Locus ID:

7249

Cytogenetics:

16p13.3

Domains:

Rap_GAP, Tuberin

Protein Families:

Druggable Genome

Protein Pathways:

Insulin signaling pathway, mTOR signaling pathway, p53 signaling pathway

MW:

40.9 kDa

Gene Summary:

Mutations in this gene lead to tuberous sclerosis complex. Its gene product is believed to be a tumor suppressor and is able to stimulate specific GTPases. The protein associates with hamartin in a cytosolic complex, possibly acting as a chaperone for hamartin. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]