

Product datasheet for **RC402632**

PKC gamma (PRKCG) (NM_002739) Human Mutant ORF Clone

Product data:

Product Type:	Mutant ORF Clones
Product Name:	PKC gamma (PRKCG) (NM_002739) Human Mutant ORF Clone
Mutation Description:	H101Y
Affected Codon#:	101
Affected NT#:	301
Nucleotide Mutation:	PRKCG Mutant (H101Y), Myc-DDK-tagged ORF clone of Homo sapiens protein kinase C, gamma (PRKCG) as transfection-ready DNA
Effect:	Spinocerebellar ataxia 14
Symbol:	PRKCG
Synonyms:	PKC-gamma; PKCC; PKCG; PKCgamma; PKCI(3); SCA14
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_002739
ORF Size:	2091 bp
Restriction Sites:	Sgfi-MluI



[View online »](#)

ORF Nucleotide
Sequence:

>RC402632 representing NM_002739
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGCTGGTCTGGGCCCGCGTAGGCGATTAGAGGGGGACCCCGGCCCTGTTTTGCAGAAAGGGG
CCCTGAGGCAGAAGGTGGTCCACGAAGTCAAGAGCCACAAGTTACCGCTCGCTTCTCAAGCAGCCAC
CTTCTGCAGCCACTGCACCGACTTCATCTGGGGTATCGAAAGCAGGGCCTGCAATGTCAAGTCTGCAGC
TTTGTGGTTTCATCGACGATGCCACGAATTTGTGACCTTCGAGTGTCCAGGCGCTGGGAAGGGCCCCCAGA
CGGACGACCCCGGAACAAATACAAGTTCGCCTGCATAGCTACAGCAGCCACCTTCTGCACCACTG
TGGCTCCCTCCTACGGGCTTGTGCACCAGGGCATGAAATGCTCCTGCTGCGAGATGAACGTGCACCGG
CGCTGTGTGCGTAGCGTGCCTCCCTGTGCGGTGTGGACCACACCGAGCGCCGGGGCGCCTGCAGCTGG
AGATCCGGGCTCCACAGCAGATGAGATCCACGTAACCTGTTGGCGAGGCCGTAACCTAATTCCTATGGA
CCCCAATGGTCTCTGATCCCTATGTGAAACTGAAGCTCATCCAGACCCTCGGAACCTGACGAAACAG
AAGACCCGAACCGTGAAAGCCACGCTAAACCCTGTGTGGAATGAGACCTTTGTGTCAACCTGAAGCCAG
GGGATGTGGAGCGCCGGCTCAGCGTGGAGGTGTGGGACTGGGACCGGACCTCCCGCAACGACTTCATGGG
GGCCATGTCTTTGGCGTCTCGGAGCTGCTCAAGGCGCCCGTGGATGGCTGGTACAAGTTACTGAACCAG
GAGGAGGGCGAGTATTACAATGTGCCGGTGGCCGATGCTGACAACTGCAGCCTCCTCCAGAAGTTTGAGG
CTTGTAACACCCCTGGAATTTGATAGAGCGGTGCGGATGGGCCCTTCTCCTCCCATCCCCCCCC
TTCCCCTAGTCCCACCGACCCCAAGCGCTGCTTCTCGGGGCGAGTCCAGGACGCTGCACATCTCCGAC
TTCAGCTTCTCATGGTCTAGGAAAAGGCAGTTTTGGAAAGGTGATGCTGGCCGAGCGCAGGGGCTCTG
ATGAGCTACGCCATCAAGATCTTAAAAAGGACGTGATCGTCCAGGACGACGATGTGGACTGCACGCT
GGTGGAGAAACGTGTGCTGGCGCTGGGGGGCCGGGGTCTTGGCGCCGGCCCACTTCTCACCCAGCTC
CACTCCACCTTCAGACCCCGGACCGCCTGATTTTCGTGATGGAGTACGTACCGGGGGAGACTTGATGT
ACCACATTCAACAGCTGGGCAAGTTTAAGGAGCCCATGCAGCGTTCTACGCGGCAGAAATCGCTATCGG
CCTCTTCTCCTTCACAATCAGGGCATCATCTACAGGGACCTGAAGCTGGACAATGTGATGCTGGATGCT
GAGGGACACATCAAGTACTGACTTTGGCATGTGTAAGGAGAACGTCTTCCCGGGACGACAACCCGCA
CCTTCTGCGGGACCCCGACTACATAGCCCGGAGATCATTGCCTACCAGCCCTATGGGAAGTCTGTGCA
TTGGTGGTCTTTGGAGTCTGCTGTATGAGATGTTGGCAGGACAGCCTCCCTTCGATGGGGAGGACGAG
GAGGAGCTGTTTCAGGCCATCATGGAACAACTGTACCTACCCCAAGTCGCTTCCCGGAAGCCGTGG
CCATCTGCAAGGGTTCTTGACCAAGCACCCAGGGAAGCGCCTGGGCTCAGGGCCTGATGGGGAACCTAC
CATCCGTGCACATGGCTTTTCCGCTGGATTGACTGGGAGCGGCTGGAACGATTGGAGATCCCGCCTCCT
TTCAGACCCCGCCGTGTGGCCGAGCGGCGAGAACTTTGACAAGTTCTTACGCGGGCGGCGCCAGCGC
TGACCCCTCCAGACCGCCTAGTCTGGCCAGCATCGACCAGGCCGATTTCAGGGCTTACCTACGTGAA
CCCCGACTTCGTGCACCCGGATGCCCGACGCCACCAGCCAGTGCCTGTGCCCGTCATG

AG**CGGACCC**ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC
TGGATTACAAGGATGACGACGA TAAGGTTTAA

Protein Sequence: >RC402632 representing NM_002739
 Red=Cloning site Green=Tags(s)

MAGLPGVGDSEGGPRPLFCRKGALRQKVVHEVKSFKFTARFFKQPTFCSHCTDFIWGIGKQGLQCQVCS
 FVVHRRCHEFVTFECPGAGKGPQTDDPRNKYFRLHSYSSPTFCDHCGSLLYGLVHQGMKSCCEMNVHR
 RCVRSVPSLCGVDHTERRGRLQLEIRAPTADEIHVTVGEARNLIPMDPNGLSDPYVKLKLIPDPRNLTKQ
 KTRTVKATLNPVWNETFVFNLKPQDVERRLSVEVWDWDRTSRNDFMGAMSFVSELLKAPVDGWYKLLNQ
 EEEGYNVPVADADNCSLLQKFEACNYPLELYERVRMGPPSSPIPSPPSPDPKRCFFGASPGRLHISD
 FSFLMVLGKGSFGKVMLAERRGSDELYAIKILKKDVIQDDDDVCTLVEKRVLALGGRGPGGRPHFLTQL
 HSTFQTPDRLYFVMEYVTGGDLMYHIQQLGKFKEPHAIFYAAEIAIGLFFLHNQGIYRDLKLDNMLDA
 EGHIKITDFGMCKENVPGTTTTRTFCGTPDYIAPEIIAYQPYGKSVDWWSFGVLLYEMLAGQPPFDGEDE
 EELFQAIMEQVTYPKSLSREAVAICKGFLTKHPGKRLGSGPDGEPTIRAHGFFRWIDWERLERLEIPPP
 FRPRPCGRSGENFDKFFTRAAPALTPPDRLLVASIDQADFQGFYVNPDFVHPDARSPTSPVVPVM

SGPTRRRLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
RefSeq:	NP_002730
RefSeq Size:	2091 bp
RefSeq ORF:	2094 bp
Locus ID:	5582
Cytogenetics:	19q13.42
Domains:	C2, pkinase, S_TK_X, TyrKc, DAG_PE-bind, S_TKc
Protein Families:	Druggable Genome, Protein Kinase
Protein Pathways:	Calcium signaling pathway, ErbB signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Gap junction, Glioma, Leukocyte transendothelial migration, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Natural killer cell mediated cytotoxicity, Non-small cell lung cancer, Pathways in cancer, Phosphatidylinositol signaling system, Tight junction, Vascular smooth muscle contraction, VEGF signaling pathway, Vibrio cholerae infection, Wnt signaling pathway
MW:	76.7 kDa

Gene Summary:

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play distinct roles in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase is expressed solely in the brain and spinal cord and its localization is restricted to neurons. It has been demonstrated that several neuronal functions, including long term potentiation (LTP) and long term depression (LTD), specifically require this kinase. Knockout studies in mice also suggest that this kinase may be involved in neuropathic pain development. Defects in this protein have been associated with neurodegenerative disorder spinocerebellar ataxia-14 (SCA14). Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015]