

Product datasheet for **RC402639**

PKC gamma (PRKCG) (NM_002739) Human Mutant ORF Clone

Product data:

Product Type:	Mutant ORF Clones
Product Name:	PKC gamma (PRKCG) (NM_002739) Human Mutant ORF Clone
Mutation Description:	Q127R
Affected Codon#:	127
Affected NT#:	380
Nucleotide Mutation:	PRKCG Mutant (Q127R), Myc-DDK-tagged ORF clone of Homo sapiens protein kinase C, gamma (PRKCG) as transfection-ready DNA
Effect:	Spinocerebellar ataxia 14
Symbol:	PRKCG
Synonyms:	PKC-gamma; PKCC; PKCG; PKCgamma; PKCI(3); SCA14
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_002739
ORF Size:	2091 bp
Restriction Sites:	SgfI-MluI



[View online »](#)

ORF Nucleotide
Sequence:

>RC402639 representing NM_002739
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCTGGTCTGGGCCCGCGTAGGCGATTAGAGGGGGACCCCGGCCCTGTTTTGCAGAAAGGGG
 CCCTGAGGCAGAAAGTGGTCCACGAAGTCAAGAGCCACAAGTTCACCGCTCGCTTCTCAAGCAGCCAC
 CTTCTGCAGCCACTGCACCGACTTCATCTGGGGTATCGAAAGCAGGGCCTGCAATGTCAAGTCTGCAGC
 TTTGTGGTTTCATCGACGATGCCACGAATTTGTGACCTTCGAGTGTCCAGGCGCTGGGAAGGGCCCCAGA
 CGGACGACCCCGGAACAACAAGTTCGCCCTGCATAGCTACAGCAGCCACCTTCTGCACCACTG
 TGGCTCCCTCCTACGGGCTTGTGCACCGGGCATGAAATGCTCCTGCTGCGAGATGAACGTGCACCGG
 CGCTGTGTGCGTAGCGTGCCTCCCTGTGCGGTGTGGACCACACCGAGCGCCGGGGCGCCTGCAGCTGG
 AGATCCGGGCTCCACAGCAGATGAGATCCACGTAACCTGTTGGCGAGGCCGTAACCTAATTCCTATGGA
 CCCCAATGGTCTCTGATCCCTATGTGAAACTGAAGCTCATCCAGACCCTCGGAACCTGACGAAACAG
 AAGACCCGAACCGTGAAGCCACGCTAAACCCTGTGTGGAATGAGACCTTTGTGTCAACCTGAAGCCAG
 GGGATGTGGAGCGCCGGCTCAGCGTGGAGGTGTGGGACTGGGACCGGACCTCCCGCAACGACTTCATGGG
 GGCCATGTCTTTGGCGTCTCGGAGTCTCAAGGCGCCCGTGGATGGCTGGTACAAGTACTGAACCAG
 GAGGAGGGCGAGTATTACAATGTGCCGGTGGCCGATGCTGACAACTGCAGCCTCCTCCAGAAGTTTGAGG
 CTTGTAACACCCCTGGAATTTGATAGAGCGGTGCGGATGGGCCCTTCTCCTCCCATCCCCCCCC
 TTCCCCTAGTCCCACCGACCCCAAGCGCTGCTTCTCGGGGCGAGTCCAGGACGCTGCACATCTCCGAC
 TTCAGCTTCTCATGGTCTAGGAAAAGGCAGTTTTGGAAAGGTGATGCTGGCCGAGCGCAGGGGCTCTG
 ATGAGCTACGCCATCAAGATCTTAAAAAGGACGTGATCGTCCAGGACGACGATGTGGACTGCACGCT
 GGTGGAGAAACGTGTGCTGGCGCTGGGGGGCCGGGGTCTTGGCGCCGGCCCACTTCTCACCCAGCTC
 CACTCCACCTTCCAGACCCCGGACCGCCTGATTTTCGTGATGGAGTACGTACCGGGGGAGACTTGATGT
 ACCACATTCAACAGCTGGGCAAGTTAAGGAGCCCATGCAGCGTTCTACGCGGCAGAAATCGCTATCGG
 CCTTCTTCTTCAATCAGGGCATCATCTACAGGGACCTGAAGCTGGACAATGTGATGCTGGATGCT
 GAGGGACACATCAAGTACTGACTTTGGCATGTGTAAGGAGAACGTCTTCCCGGGACGACAACCCGCA
 CCTTCTGCGGGACCCCGACTACATAGCCCGGAGATCATTGCCTACCAGCCCTATGGGAAGTCTGTGCA
 TTGGTGGTCTTTGGAGTCTGCTGTATGAGATGTTGGCAGGACAGCCTCCCTTCGATGGGGAGGACGAG
 GAGGAGCTGTTTCAGGCCATCATGGAACAACTGTACCTACCCCAAGTCGCTTCCCGGAAGCCGTGG
 CCATCTGCAAGGGTTCTTGACCAAGCACCCAGGGAAGCGCCTGGGCTCAGGGCCTGATGGGGAACCTAC
 CATCCGTGCACATGGCTTTTCCGCTGGATTGACTGGGAGCGGCTGGAACGATTGGAGATCCCGCCTCCT
 TTCAGACCCCGCCGTGTGGCCGACGGCGGAGAACTTTGACAAGTCTTACGCGGGCGGCGCCAGCGC
 TGACCCCTCCAGACCGCTAGTCTGGCCAGCATCGACCAGGCCGATTTCAGGGCTTACCTACGTGAA
 CCCCAGCTTCGTGCACCCGGATGCCCGACGCCACCAGCCAGTGCCTGTGCCCGTCATG

AG**CGGACCC**ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC
 TGGATTACAAGGATGACGACGA TAAGGTTTAA

Protein Sequence: >RC402639 representing NM_002739
 Red=Cloning site Green=Tags(s)


MAGLPGVGDSEGGPRPLFCRKALRQKVVHEVKSFKFTARFFKQPTFCSHCTDFIWGIGKQGLQCQVCS
 FVVHRRCHEFVTFECPGAGKGPQTDDPRNKKHFRHLSYSSPTFCDHCGSLLYGLVHRGMKSCCEMNVHR
 RCVRSVPSLGGVDHTERRGRLQLEIRAPTADEIHVTVGEARNLIPMDPNGLSDPYVKLKLIPDPRNLTKQ
 KTRTVKATLNPVWNETFVFNLKPGDVERRLSVEVWDWDRTSRNDFMGAMSFVSELLKAPVDGWYKLLNQ
 EEEGYNVPVADADNCSLLQKFEACNYPLELYERVRMGPSSSPIPSPSPSPTDPKRCFFGASPGRLHISD
 FSFLMVLGKGSFGKVMMLAERGSDELYAIKILKKDVIQDDDDVCTLVEKRVLALGGRGPGGRPHFLTQL
 HSTFQTPDRLYFVMEYVTGGDLMYHIQQLGKFKEPHAAFYAAEIAIGLFFLHNQGIYRDLKLDNMLDA
 EGHIKITDFGMCKENVPGTTTTRTFCGTPDYIAPEIIAYQPYGKSVDDWWSFGVLLYEMLAGQPPFDGEDE
 EELFQAIMEQTVTPKSLSREAVAICKGFLTKHPGKRLGSGPDGEPTIRAHGFFRWIDWERLERLEIPPP
 FRPRPCGRSGENFDKFFTRAAPALTPPDRLLASIDQADFQGFYVNPDFVHPDARSPTSPVVPVM

SGPTRRRLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
RefSeq:	NP_002730
RefSeq Size:	2091 bp
RefSeq ORF:	2094 bp
Locus ID:	5582
Cytogenetics:	19q13.42
Domains:	C2, pkinase, S_TK_X, TyrKc, DAG_PE-bind, S_TKc
Protein Families:	Druggable Genome, Protein Kinase
Protein Pathways:	Calcium signaling pathway, ErbB signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Gap junction, Glioma, Leukocyte transendothelial migration, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Natural killer cell mediated cytotoxicity, Non-small cell lung cancer, Pathways in cancer, Phosphatidylinositol signaling system, Tight junction, Vascular smooth muscle contraction, VEGF signaling pathway, Vibrio cholerae infection, Wnt signaling pathway
MW:	76.7 kDa

Gene Summary:

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play distinct roles in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase is expressed solely in the brain and spinal cord and its localization is restricted to neurons. It has been demonstrated that several neuronal functions, including long term potentiation (LTP) and long term depression (LTD), specifically require this kinase. Knockout studies in mice also suggest that this kinase may be involved in neuropathic pain development. Defects in this protein have been associated with neurodegenerative disorder spinocerebellar ataxia-14 (SCA14). Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015]