

Product datasheet for **RC402983**

BRCA1 (NM_007294) Human Mutant ORF Clone

Product data:

Product Type:	Mutant ORF Clones
Product Name:	BRCA1 (NM_007294) Human Mutant ORF Clone
Mutation Description:	S868X
Affected Codon#:	868
Affected NT#:	2603
Nucleotide Mutation:	BRCA1 Mutant (S868X), Myc-DDK-tagged ORF clone of Homo sapiens breast Cancer, early onset (BRCA1), transcript variant 1 as transfection-ready DNA
Effect:	Breast cancer
Symbol:	BRCA1
Synonyms:	BRCAI; BRCC1; BROVCA1; FANCS; IRIS; PNCA4; PPP1R53; PSCP; RNF53
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_007294
ORF Size:	2601 bp
Restriction Sites:	Sgfl-Mlul



[View online »](#)

**ORF Nucleotide
Sequence:**

>RC402983 representing NM_007294
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGATTTATCTGCTCTTCGCGTTGAAGAAGTACAAAATGTCATTAATGCTATGCAGAAAATCTTAGAGT
 GTCCCATCTGTCTGGAGTTGATCAAGGAACCTGTCTCCACAAAGTGTGACCACATATTTTGCAAATTTG
 CATGCTGAAACTTCTCAACCAGAAGAAAGGGCCTTACAGTGTCTTTATGTAAGAATGATATAACCAAA
 AGGAGCCTACAAGAAAGTACGAGATTTAGTCAACTTGTGAAGAGCTATTGAAAAATTTGTGCTTTTC
 AGCTTGACACAGGTTTGGAGTATGCAAACAGCTATAATTTTGCAAAAAGGAAAAAATACTCTCTGAACA
 TCTAAAAGATGAAGTTTCTATCATCCAAAGTATGGGCTACAGAAAACCGTCCAAAAGACTTCTACAGAGT
 GAACCCGAAAAATCCTTCCTTGCAGGAAACAGTCTCAGTGTCCAACCTCTAACCTTGGAACTGTGAGAA
 CTCTGAGGACAAAGCAGCGGATACAACCTCAAAAGACGTCTGTCTACATTGAATTGGGATCTGATTCTTC
 TGAAGATACCGTTAATAAGGCAACTTATTGCAGTGTGGGAGATCAAGAATTGTTACAAAATCACCCCTCAA
 GGAACCCAGGGATGAAATCAGTTTGGATTCTGCAAAAAAGGCTGCTTGTGAATTTTCTGAGACGGATGAA
 CAAATACTGAACATCATCAACCCAGTAATAATGATTTGAACACCACTGAGAAGCGTGCAGCTGAGAGGCA
 TCCAGAAAAGTATCAGGGTAGTTCTGTTTCAAACCTGTCATGTGGAGCCATGTGGCACAAAATACTCATGCC
 AGCTCATTACAGCATGAGAACAGCAGTTTATTACTCACTAAAGACAGAAATGAATGTAGAAAAGGCTGAAT
 TCTGTAATAAAAAGCAAACAGCCTGGCTTAGCAAGGAGCCAACATAACAGATGGGCTGGAAGTAAGGAAAC
 ATGTAATGATAGGCGGACTCCAGCACAGAAAAAAGGTAGATCTGAATGCTGATCCCTGTGTGAGAGA
 AAAGAAATGGAATAAGCAGAACTGCCATGCTCAGAGAATCCTAGAGATACTGAAGATGTTCCCTGGATAA
 CACTAAATAGCAGCATTAGAAAAGTTAATGAGTGGTTTTCCAGAAGTATGAACCTGTTAGTTCTGATGA
 CTCACATGATGGGGAGTCTGAATCAAATGCCAAAGTAGCTGATGTATTGGACGTTCTAAATGAGGTAGAT
 GAATATTCTGGTTCTTACAGAAAAATAGACTTACTGGCCAGTGATCCTCATGAGGCTTTAATATGAAAA
 GTGAAAGAGTTCACTCCAAATCAGTAGAGAGTAATATTGAAGACAAAATATTTGGGAAAACCTATCGGAA
 GAAGGCAAGCCTCCCAACTTAAGCCATGTAACGAAAAATCTAATTATAGGAGCATTGTTACTGAGCCA
 CAGATAATACAAGAGCGTCCCTCACAAATAAATTAAGCGTAAAAGGAGACCTACATCAGGCTTCATC
 CTGAGGATTTTATCAAGAAAGCAGATTTGGCAGTTCAAAGACTCCTGAAATGATAAATCAGGAACTAA
 CCAAACGAGCAGAATGGTCAAGTATGAATATTACTAATAGTGGTCATGAGAATAAAACAAAAGGTGAT
 TCTATTCAGAATGAGAAAAATCCTAACCAATAGAATCACTCGAAAAAGAAATCTGCTTTCAAACGAAAG
 CTGAACCTATAAGCAGCAGTATAAGCAATATGGAACCTCGAATTAATATCCACAATTCAAAAGCACCTAA
 AAAGAATAGGCTGAGGAGGAAGTCTTCTACCAGGCATATTCATGCGCTTGAACCTAGTAGTCAGTAGAAAT
 CTAAGCCACCTAATTGTACTGAATTGCAAATGATAGTTGTTCTAGCAGTGAAGAGATAAAGAAAAAAA
 AGTACAACCAATGCCAGTCAGGCACAGCAGAAACCTACAACCTCATGGAAGGTAAGAACCTGCAACTGG
 AGCCAAGAAGAGTAACAAGCCAAATGAACAGACAAGTAAAAGACATGACAGCGTACTTTCCAGAGCTG
 AAGTTAACAAAATGCACCTGGTTCTTTTACTAAGTGTCAAATACCAGTGAACCTAAAGAATTTGTCAATC
 CTAGCCTTCCAAGAGAAGAAAAAGAAGAGAACTAGAAACAGTTAAAGTGTCTAATAATGCTGAAGACCC
 CAAAGATCTCATGTTAAGTGGAGAAAGGTTTTGCAAACCTGAAAGATCTGTAGAGAGTAGCAGTATTTCA
 TTGGTACTGGTACTGATTATGGCACTCAGGAAAGTATCTCGTTACTGGAAGTTAGCACTTAGGGAAAGG
 CAAAAACAGAACCAAAATAAATGTGTGAGTCAAGTGTGCAGCATTGAAAACCCCAAGGACTAATTCATGG
 TTGTTCCAAAGATAATAGAAATGACACAGAAGGCTTTAAGTATCCATTGGGACATGAAGTTAACCACAGT
 CGGAAACAAGCATAGAAATGGAAGAAAGTGAACCTTGATGCTCAGTATTTGCAGAATACATTCAAGGTTT
 CAAAGCGCCAG

AG**CGGACCG**ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC
 TGGATTACAAGGATGACGACGA TAAGGTTTAA

Protein Sequence: >RC402983 representing NM_007294
 Red=Cloning site Green=Tags(s)



MDLSALRVEEVQNVINAMQKILECPICLLELIKEPVSTKCDHIFCKFCMLKLLNQKKGPSQCPLCKNDITK
 RSLQESTRFSQLVEELLKIICAFQLDTGLEAYANSYNFAKKENNSPEHLKDEVSI IQSMGYRNRARLLQS
 EPENPSLQETSLSVQLSNLGTVRTLRKQRIQPQKTSVYIELGSDSSEDTVNKATYCSVGDQELLQITPQ
 GTRDEISLDSAKKAACEFSETDVTNTEHHQPSNNDLNTTEKRAAERHPEKYQGSSVSNLHVPCGTNTHA
 SSLQHENSLLLLTKDRMNVKEAEFCNKSKQPGLARSQHNRWAGSKETCNDRRTPSTEKKVDLNADPLCER
 KEWKNQKLPCCSENPRDTEVPWITLNSSIQKVNEWFSRSEDELLGSDSDHGESESNKAVDVLVDLNEVD
 EYSGSSEKIDLLASDPHEALICKSERVHKSVESNIEDKIFGKTYRKKASLPNLSHVTENLIIGAFVTEP
 QIIQERPLTNKLRKRRTSGLHPEDFIKKADLAVQKTPEMINQGTNQTEQNGQVMNITNSGHENKTKGD
 SIQNEKNPNPIESLEKESAFKTKAEPISSSISNMELELNIHNSKAPKKNRLRRKSSTRHIALELVVSRN
 LSPNCTELQIDSCSSSEI KKKKYNQMPVRRSRLQLMEGKEPATGAKKSNKPNEQTSKRHDSDTFPEL
 KLTNAPGSFTKCSNTSELKEFVNPSLPREEKEELETVKVSNNAEDPKDMLSGERVLQTERSVESSSIS
 LVPGTGYGTQESISLLEVSTLGAKTEPNKCVSQCAAFENPKGLIHGCSKDNRRNDEGFKYPLGHEVNHS
 RETSIEMEESELDAQYLQNTFKVSKRQ

SGP TRTRRL EQKL I SEEDLA AND I LDYK DDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:



* The last codon before the Stop codon of the ORF

OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NP_009225
RefSeq Size:	2601 bp
RefSeq ORF:	5592 bp
Locus ID:	672
Cytogenetics:	17q21.31
Domains:	BRCT, RING
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Ubiquitin mediated proteolysis
MW:	95.4 kDa

Gene Summary:

This gene encodes a 190 kD nuclear phosphoprotein that plays a role in maintaining genomic stability, and it also acts as a tumor suppressor. The BRCA1 gene contains 22 exons spanning about 110 kb of DNA. The encoded protein combines with other tumor suppressors, DNA damage sensors, and signal transducers to form a large multi-subunit protein complex known as the BRCA1-associated genome surveillance complex (BASC). This gene product associates with RNA polymerase II, and through the C-terminal domain, also interacts with histone deacetylase complexes. This protein thus plays a role in transcription, DNA repair of double-stranded breaks, and recombination. Mutations in this gene are responsible for approximately 40% of inherited breast cancers and more than 80% of inherited breast and ovarian cancers. Alternative splicing plays a role in modulating the subcellular localization and physiological function of this gene. Many alternatively spliced transcript variants, some of which are disease-associated mutations, have been described for this gene, but the full-length natures of only some of these variants has been described. A related pseudogene, which is also located on chromosome 17, has been identified. [provided by RefSeq, May 2020]