

## Product datasheet for **RC403506**

### MEK2 (MAP2K2) (NM\_030662) Human Mutant ORF Clone

#### Product data:

Product Type:	Mutant ORF Clones
Product Name:	MEK2 (MAP2K2) (NM_030662) Human Mutant ORF Clone
Mutation Description:	Y134C
Affected Codon#:	134
Affected NT#:	401
Nucleotide Mutation:	MAP2K2 Mutant (Y134C), Myc-DDK-tagged ORF clone of Homo sapiens mitogen-activated protein kinase kinase 2 (MAP2K2) as transfection-ready DNA
Effect:	Crdio-fio-uneous syndrome
Symbol:	MAP2K2
Synonyms:	CFC4; MAPKK2; MEK2; MKK2; PRKMK2
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_030662
ORF Size:	1200 bp
Restriction Sites:	Sgfi-Mlul



[View online »](#)

**ORF Nucleotide Sequence:**

>RC403506 representing NM\_030662  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**GCGATCGCC**

ATGCTGGCCCGGAGGAAGCCGGTGTGCCGGCGCTCACCATCAACCCTACCATCGCCGAGGGCCCATCCC  
 CTACCAGCGAGGGCGCCTCCGAGGCAAACCTGGTGGACCTGCAGAAGAAGCTGGAGGAGCTGGAACCTTGA  
 CGAGCAGCAGAAGAAGCGGCTGGAAGCCTTTCTCACCCAGAAAGCCAAGGTGGCGAACTCAAAGACGAT  
 GACTTCGAAAGGATCTCAGAGCTGGGCGCGGGCAACGGCGGGGTGGTCACCAAAGTCCAGCACAGACCCT  
 CGGGCCTCATCATGGCCAGGAAGCTGATCCACCTTGAGATCAAGCCGGCCATCCGGAACCAGATCATCCG  
 CGAGCTGCAGGTCTGCACGAATGCAACTCGCCGTACATCGTGGGCTTCTGCGGGCCTTCTACAGTGAC  
 GGGGAGATCAGCATTTGCATGGAACACATGGACGGCGGCTCCCTGGACCAGGTGCTGAAAGAGGCCAAGA  
 GGATCCCGAGGAGATCTGGGAAAGTCAAGTCCGCGTTCTCCGGGGCTTGGCGTACCTCCGAGAGAA  
 GCACCAGATCATGCACCGAGATGTGAAGCCCTCAACATCCTCGTGAAGTCTAGAGGGGAGATCAAGCTG  
 TGTGACTTCGGGGTGAAGCGCCAGCTCATCGACTCCATGGCCAACTCCTTCGTGGGCACGCGCTCCTACA  
 TGGCTCCGGAGCGGTTGACAGGGCACACATTACTCGGTGCAGTCCGACATCTGGAGCATGGGCTGTCCCT  
 GGTGGAGCTGGCCGTCGGAAGGTACCCCATCCCCCGCCGACGCCAAAGAGCTGGAGGCCATCTTTGGC  
 CGGCCCGTGGTCGACGGGAAGAAGGAGAGCCTCACAGCATCTCGCCTCGGCCGAGGCCCGCCGCGCC  
 CCGTCAGCGGTACGGGATGGATAGCCGGCCTGCCATGGCCATCTTTGAACTCTGGACTATATTGTGAA  
 CGAGCCACCTCCTAAGCTGCCAACGGTGTGTTACCCCCGACTTCCAGGAGTTGTCAATAAATGCCTC  
 ATCAAGAACCCAGCGGAGCGGGCGGACCTGAAGATGCTCACAACCCACACCTTCAAGCGGTCCGAGG  
 TGAAGAAGTGGATTTTCCCGGCTGTTGTGTAAAACCTGCGGCTGAACCAGCCCGGCACACCCACGCG  
 CACCGCGTG

AG**GCGACCG**ACGCGTACGCGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC  
 TGGATTACAAGGATGACGACGA TAAGGTTTAA

**Protein Sequence:**


>RC403506 representing NM\_030662  
 Red=Cloning site Green=Tags(s)

MLARRKPVLPAL TINPTIAEGPSPTSEGASEANLVDLQKKLEELDEQKKRLEAFLTQKAKVGELKDD  
 DFERISELGAGNGGVTKVQHRPSGLIMARKLIHLEIKPAIRNQIIRELQVLHECNSPYIVGFCGAFYSD  
 GEISICMEHMDGGSLDQVLKEAKRIPEEILGKVSIAVLRGLAYLREKHQIMHRDVKPSNILVNSRGEIKL  
 CDFGVSGQLIDSMANSFVGRSYMAPERLQGTHYSVQSDIWSMGLSLVELAVGRYPPIPPDAKELEAIFG  
 RPVVDGEEGEPHSISPRPRPPGRPVSGHGMSRPAMAI FELLDYIVNEPPPPLPNGVFTPDFQEFVNKCL  
 IKNPAERADLKMLTNHTFIKRSEVEEVDFAWGLCKTLRLNQPPTPTAV

**SGP**TRRRLEQKLI SEEDLAANDILDYKDDDDKV

**Restriction Sites:**

Sgfl-Mlul

**Cloning Scheme:**

**OTI Disclaimer:**

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:**

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:**

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**RefSeq:**

[NP\\_109587](#)

**RefSeq Size:**

1200 bp

**RefSeq ORF:**

1203 bp

**Locus ID:**

5605

**Cytogenetics:**

19p13.3

**Domains:**

pkinase, TyrKc, S\_TKc

**Protein Families:**

Druggable Genome, Protein Kinase

**Protein Pathways:**

Acute myeloid leukemia, B cell receptor signaling pathway, Bladder cancer, Chronic myeloid leukemia, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Gap junction, Glioma, GnRH signaling pathway, Insulin signaling pathway, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Melanoma, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Non-small cell lung cancer, Pathways in cancer, Prion diseases, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway, Thyroid cancer, Toll-like receptor signaling pathway, Vascular smooth muscle contraction, VEGF signaling pathway

**MW:** 44 kDa

**Gene Summary:** The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, cognitive disability, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene. [provided by RefSeq, Jul 2008]