

## Product datasheet for **RG200398**

### FHL2 (NM\_201557) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	FHL2 (NM_201557) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	FHL2
Synonyms:	AAG11; DRAL; FHL-2; SLIM-3; SLIM3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG200398 representing NM_201557 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGACTGAGCGCTTTGACTGCCACCATTGCAACGAATCTCTTTGGCAAGAAGTACATCCTGCGGGAGG  
AGAGCCCCTACTGCGTGGTGTGCTTTGAGACCCTGTTCCGCAACACCTGCGAGGAGTGTGGGAAGCCCAT  
CGGCTGTGACTGCAAGGACTTGTCTTACAAGGACCGGCACTGGCATGAAGCCTGTTCCACTGCTCGCAG  
TGCAGAACTCACTGGTGGACAAGCCCTTGTGCCAAGGAGGACCAGCTGCTCTGTACAGACTGCTATT  
CCAACGAGTACTCATCCAAGTGCCAGGAATGCAAGAAGACCATCATGCCAGGTACCGCAAGATGGAGTA  
CAAGGGCAGCAGCTGGCATGAGACCTGCTTCATCTGCCACCGCTGCCAGCAGCCAATTGGAACCAAGAGT  
TTCATCCCCAAAGACAATCAGAATTTCTGTGTGCCCTGCTATGAGAAAACAACATGCCATGCAGTGCCTTC  
AGTGCAAAAAGCCCATCACACGGGAGGGGTCACTTACCGGGAGCAGCCCTGGCACAAGGAGTGTTCGT  
GTGCACCGCCTGCAGGAAGCAGCTGTCTGGCAGCGCTTACAGCTCGCGATGACTTTGCCTACTGCCTG  
AACTGCTTCTGTGACTTGTATGCCAAGAAGTGTGCTGGGTGCACCAACCCCATCAGCGGACTTGGTGGCA  
CAAATACATCTCCTTTGAGGAACGGCAGTGGCATAACGACTGCTTTAACTGTAAGAAGTGTCCCTCTC  
ACTGGTGGGGCGTGGCTTCTCACAGAGAGGGACGACATCCTGTGCCCGACTGTGGAAAGACATC

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG200398 representing NM\_201557  
 Red=Cloning site Green=Tags(s)

MTERFDCHHCNESLFGKKYILREESPYCVVCFETLFANTCEECKPIGCDCKDLSYKDRHWHEACFHCSQ  
 CRNSLVDKPFPAKEDQLLCTDCYSNEYSSKQCECKKTIIMPGTRKMEYKGSWHETCFICHRCQQPIGTKS  
 FIPKDNQNFVPCYEKQHAMQCVQCKKPIITGGVTYREQPWHKECFVCTACRQKLSGQRFTARDDFAYCL  
 NCFCDLYAKKCAGCTNPIISGLGGTKYISFEERQWHNDCFNCKKCSLSLVGRGFLTERDDILCPDCGKDI

TRTRPLE - GFP Tag - V


**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shutting:



**Plasmid Map:**



**ACCN:** NM\_201557

**ORF Size:** 837 bp

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_201557.4</a>
<b>RefSeq Size:</b>	1424 bp
<b>RefSeq ORF:</b>	840 bp
<b>Locus ID:</b>	2274
<b>UniProt ID:</b>	<a href="#">Q14192</a>
<b>Cytogenetics:</b>	2q12.2
<b>Protein Families:</b>	Druggable Genome
<b>Gene Summary:</b>	This gene encodes a member of the four-and-a-half-LIM-only protein family. Family members contain two highly conserved, tandemly arranged, zinc finger domains with four highly conserved cysteines binding a zinc atom in each zinc finger. This protein is thought to have a role in the assembly of extracellular membranes. Also, this gene is down-regulated during transformation of normal myoblasts to rhabdomyosarcoma cells and the encoded protein may function as a link between presenilin-2 and an intracellular signaling pathway. Multiple alternatively spliced variants encoding different isoforms have been identified. [provided by RefSeq, Jan 2016]