

Product datasheet for **RG200436**

PSMD8 (NM_002812) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PSMD8 (NM_002812) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	PSMD8
Synonyms:	HEL-S-91n; HIP6; HYPF; Nin1p; p31; Rpn12; S14
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG200436 representing NM_002812 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGTACGAGCAACTCAAGGGCGAGTGAACCGTAAAAGCCCCAATCTTAGCAAGTGCAGGGAAGAGCTGG
GTCGACTCAAGCTAGTTCTTCTGGAGCTCAACTTCTTGCCAACACAGGGACCAAGCTGACCAAACAGCA
GCTAATTCTGGCCGTGACATACTGGAGATCGGGGCCAATGGAGCATCCTACGCAAGGACATCCCCTCC
TTCGAGCGCTACATGGCCAGCTCAAATGCTACTACTTTGATTACAAGGAGCAGCTCCCCGAGTCAGCCT
ATATGCACCAGCTCTTGGCCTCAACCTCCTCTCCTGCTGTCCAGAACCGGTGGCTGAGTTCACAC
GGAGTTGGAGCGGCTGCCTGCCAAGGACATACAGACCAATGTCTACATCAAGCACCCAGTGTCCCTGGAG
CAATACCTGATGGAGGGCAGCTACAACAAAGTGTCTCCTGGCCAAGGGTAACATCCCCGCCGAGAGCTACA
CCTTCTTCATTGACATCCTGCTCGACACTATCAGGGATGAGATCGCTGGGTGCATCGAGAAGGCCTACGA
GAAAATCCTTTTCACTGAGGCCACCCGGATCCTCTTCTTCAACACACCCAAAAAGATGACAGACTACGCC
AAGAAGCGAGGGTGGTCTCCTGGGCCCAACAATACTACAGTTTTGCCAGCCAGCAGCAGAAGCCGGAAG
ACACCACCATTCCCTCCACAGAAGTGGCCAAACAGGTCATCGAGTATGCCCGGCAGCTGGAGATGATCGT
C

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG200436 representing NM_002812
 Red=Cloning site Green=Tags(s)

MYEQLKGEWNRKSPNLSKCGEELGRLKLVLELNFLPTTGKLTQQQLILARDILEIGAQWSILRKDIPS
 FERYMAQLKCYFDYKEQLPESAYMHQLLGLNLLFLLSQNRVAEFHTELERLPAKDIQTNVYIKHPVSLE
 QYLMEGSYNKVFLAKGNIPAESYTFIDILLDIRDEIAGCIEKAYEKILFTEATRILFFNTPKKMTDYA
 KKRGWVLGPNNYSFASQQKPEDTTIPSTELAKQVIEYARQLEMIV

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_002812

ORF Size: 771 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_002812.3](#), [NP_002803.1](#)

RefSeq Size: 1556 bp

RefSeq ORF: 1053 bp

Locus ID: 5714

UniProt ID: [P48556](#)


Cytogenetics: 19q13.2

Domains: Nin1_C

Protein Pathways: Proteasome

Gene Summary: The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. A pseudogene has been identified on chromosome 1. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RG200436