

Product datasheet for **RG200484**

CPT2 (NM_000098) Human Tagged ORF Clone

Product data:


Product Type:	Expression Plasmids
Product Name:	CPT2 (NM_000098) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	CPT2
Synonyms:	CPT1; CPTASE; IIAE4
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



[View online »](#)

Plasmid Map:



ACCN: NM_000098

ORF Size: 1974 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_000098.1](#), [NP_000089.1](#)

RefSeq Size:	3090 bp
RefSeq ORF:	1977 bp
Locus ID:	1376
UniProt ID:	P23786
Cytogenetics:	1p32.3
Domains:	Carn_acyltransf
Protein Families:	Druggable Genome
Protein Pathways:	Fatty acid metabolism, PPAR signaling pathway
Gene Summary:	The protein encoded by this gene is a nuclear protein which is transported to the mitochondrial inner membrane. Together with carnitine palmitoyltransferase I, the encoded protein oxidizes long-chain fatty acids in the mitochondria. Defects in this gene are associated with mitochondrial long-chain fatty-acid (LCFA) oxidation disorders. [provided by RefSeq, Jul 2008]