

Product datasheet for **RG200495**

CDK1 (NM_001786) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CDK1 (NM_001786) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	CDK1
Synonyms:	CDC2; CDC28A; P34CDC2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG200495 representing NM_001786 Red=Cloning site Blue=ORF Green=Tags(s)

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGAAGATTATACCAAAATAGAGAAAATTGGAGAAGGTACCTATGGAGTTGTGTATAAGGGTAGACACA
AAACTACAGGTCAAGTGGTAGCCATGAAAAAATCAGACTAGAAAGTGAAGAGGAAGGGTTCCTAGTAC
TGCAATTCGGGAAATTTCTCTATTAAAGGAACTTCGTCATCCAAATATAGTCAGTCTTCAGGATGTGCTT
ATGCAGGATTCCAGGTTATATCTCATCTTTGAGTTTCTTCCATGGATCTGAAGAAATACTTGGATTCTA
TCCCTCCTGGTCAGTACATGGATTCTTCACTTGTAAAGAGTTATTTATACCAATCCTACAGGGGATTGT
GTTTTGTCACTCTAGAAGAGTTCTTACAGAGACTTAAACCTCAAAATCTTTGATTGATGACAAAGGA
ACAATTAAGTGGCTGATTTTGGCCTTGCCAGAGCTTTTGAATACCTATCAGAGTATATACACATGAGG
TAGTAACACTCTGGTACAGATCTCCAGAAGTATTGCTGGGGTCAGCTCGTTACTCAACTCCAGTTGACAT
TTGGAGTATAGGCACCATATTTGCTGAAGTACCACTAAGAAACCACTTTTCCATGGGGATTGAGAAATT
GATCAACTCTTCAGGATTTTCAGAGCTTTGGGCACTCCCAATAATGAAGTGTGCCAGAAGTGAATCTT
TACAGGACTATAAGAATACATTTCCCAATGGAACCAAGCAAGCTAGCATCCCATGTCAAAACTTGGA
TGAAAATGGCTTGGATTTGCTCTCGAAAATGTTAATCTATGATCCAGCCAAACGAATTTCTGGCAAAATG
GCACTGAATCATCCATATTTAATGATTGGACAATCAGATTAAGAAGATG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG200495 representing NM_001786
 Red=Cloning site Green=Tags(s)

MEDYTKIEKIGEGTYGVVYKGRHKTTGQVAMKKIRLESEEEGVPSTAIREISLLKELRHPNIVSLQDVL
 MQDSRLYLIFEFLSMDLKKYLDIIPPGQYMDSSLVKSILYQILQGI V FCHSRRVLHRDLKPQNLLIDDKG
 TIKLADFGLARAFGIPIRVYTHEVVT LWYRSPEVLLGSARYSTPVDIWSIGTIFAELATKKPLFHGDSEI
 DQLFRIFRALGTPNNEVWPEVESLQDYKNTFPKWKPGSLASHVKNLDENGLDLLSKMLIYDPAKRISGKM
 ALNHPYFNDLDNQIKKM

TRTRPLE – GFP Tag – V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001786

ORF Size: 891 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001786.5</u>
RefSeq Size:	1235 bp
RefSeq ORF:	894 bp
Locus ID:	983
UniProt ID:	<u>P06493</u>
Cytogenetics:	10q21.2
Domains:	pkinase, TyrKc, S_TKc
Protein Families:	Druggable Genome, Protein Kinase, Stem cell - Pluripotency
Protein Pathways:	Cell cycle, Gap junction, Oocyte meiosis, p53 signaling pathway, Progesterone-mediated oocyte maturation
Gene Summary:	<p>The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009]</p>

Product images:



Circular map for RG200495