

Product datasheet for **RG200523**

HSPA2 (NM_021979) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	HSPA2 (NM_021979) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	HSPA2
Synonyms:	HSP70-2; HSP70-3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RG200523 representing NM_021979
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGCATCGCC

ATGTCTGCCCGTGGCCCGGCTATCGGCATCGACCTGGGCACCACCTATTCGTGCGTCGGGGTCTTCCAAC
 ATGGCAAGGTGGAGATCATCGCCAACGACCAGGGCAATCGCACCACCCAGCTACGTGGCCTTCACGGA
 CACCGAGCGCCTCATCGCGCAGCGCCCAAGAACCAGGTGGCCATGAACCCACCAACACCATCTTCGAC
 GCCAAGAGGCTGATTGGACGAAATTCGAGGATGCCACAGTGCAGTCGGATATGAAACACTGGCCGTTC
 GGGTGGTGGAGGAGGCAAGCCAAAGTGAAGTAGAGTACAAGGGGAGACCAAGACCTTCTTCCC
 AGAGGAGATATCCTCCATGGTCTCACGAAGATGAAGGAGATCGCGGAAGCCTACCTGGGGGCAAGGTG
 CACAGCGCGGTCATAACGGTCCCGGCTATTTCAACGACTCGCAGCGCCAGGCCACCAAGGACGCAGGCA
 CCATCACGGGGTCAATGTGCTGCGCATCATCAACGAGCCACGGCGGCCCATCGCCTACGGCCTGGA
 CAAGAAGGGTGGCGGGCGGCGAGAAGAAGTGTCTCATCTTTGACCTGGGCGGTGGCACTTTCGACGTG
 TCCATCTGACCATCGAGGATGGCATCTTCGAGGTGAAGTCCACGGCCGGCGACACCCACCTGGGCGGTG
 AGGACTTCGACAACCGCATGGTGAAGCCACTGGCGGAGGAGTTCAAGCGCAAGCACAAGAAGGACATTGG
 GCCCAACAAGCGCGCCTGAGGGGCTGCGCACCGCTTGCAGCGCGCCAAGCGCACCTGAGCTCGTCC
 ACGCAGGCGAGCATCGAGATCGACTCGCTACGAGGGCGTGGACTTCTATACGTCCATCACGCGCGCC
 GCTTCGAGGAGCTCAATGCCGACCTTTTCGCGGGACCTGGAGCCGGTGGAGAAGGCGCTGCGCGACGC
 CAAGCTGGACAAGGGCCAGATCCAGGAGATCGTGTGGTGGGCGGCTCCACTCGTATCCCAAGATCCAG
 AAGCTGCTGCAGATTTCTTCAACGGCAAGGAGCTGAACAAGAGCATCAACCCGACGAGGCGGTGGCCT
 ATGGCGCCGCGGTGCAGGCGGCCATCTCATCGCGCACAATCAGAGAATGTGACGAGACCTGCTGCTACT
 CGACGTGACCCCGTTGTGCTGGGCATCGAGACAGCTGGCGGTGTCATGACCCCACTCATCAAGAGGAAC
 ACCACGATCCCCACCAAGCAGACGACGACCTTACCACCTACTCGGACAACCAGAGCAGCGTACTGGTGC
 AGGTATACGAGGGCGAACGGCCATGACCAAGGACAATAACCTGTGGGCAAGTTCGACCTGACCGGGAT
 TCCCCCTGCGCCTCGCGGGTCCCCAAATCGAGGTTACCTTCGACATTGACGCAATGGCATCCTTAAC
 GTTACCGCGCGACAAGAGCACCGGTAAAGAAAAAAAATCACCATACCAATGACAAAGGTGCTCTGA
 GCAAGGACGACATTGACCGGATGGTGCAGGAGGCGGAGCGGTACAATCGGAAGATGAGGCGAATCGCGA
 CCGAGTCGCGGCAAAAACGCCCTGGAGTCTATACTACAACATCAAGCAGACGGTGAAGACGAGAAA
 CTGAGGGCAAGATTAGCGAGCAGGACAAAACAAGATCTCGACAAGTGTGAGGAGGTGATCAACTGGC
 TCGACCGAAACCAGATGGCAGAGAAAGATGAGTATGAACAACAAGCAGAAAGAGCTCGAAAGATTTGCAA
 CCCCATCATCAGCAAACCTTACCAAGGTGGTCTGGCGGCGCAGCGCGCGGCGGTTACGAGACCTCC
 GGGGACCCACCATCGAAGAAGTGGAC

ACGCGTACGCGGCGGCTCGAG – GFP Tag – GTTTAA

Protein Sequence:

>RG200523 representing NM_021979
 Red=Cloning site Green=Tags(s)

MSARGPAIGIDLTTYSCVGVFQHGKVEI IANDQGNRTTPSYVAFDTERLIGDAAKNQVAMNPTNTIFD
 AKRLIGRKFEDATVQSDMKHWPFRVSEGGPKVQVEYKGETKTFPPEI SSMVLTKMKEIAEAYLGKV
 HSAVITVPAYFNDSQRQATKDAGTITGLNVLRIINEPTAAAIAYGLDKKGCAGGEKNVIFDLGGGTFDV
 SILTIEDGIFEVKSTAGDTHLGGEDFDNRMVSHLAEFEKRRKHKDIPGNKRAVRRRLTACERAKRTLSS
 TQASIEIDSLYEGVDFYTSITRARFEELNADLFRGTLEPVEKALRDAKLDKGQIQEIVLVGGSTRIPKIQ
 KLLQDFNFKELNKSINPDEAVAYGAAVQAAILIGDKSENVQDLLLDVTPLSLGIETAGVMTPLIKRN
 TTIPKQTQFTTYSQSSVLVQVYEGERAMTKDNNLLGKFDLGTGIPAPRGVPIEVTFDIDANGILN
 VTAADKSTGKENKITITNDKGRLSKDDIDRMVQEAERYKSEDEANRDRVAAKNALESYTYNIKQTVDEK
 LRGKISEQDNKILDKCQEVINWLDNRNQMAEKDEYEHKQKELERVCNPIISKLYQGGPGGGGGGGSGS
 GGPTIEEVD

TRTRPLE – GFP Tag – V

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:



ACCN: NM_021979

ORF Size: 1917 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_021979.3
RefSeq Size:	2496 bp
RefSeq ORF:	1920 bp
Locus ID:	3306
UniProt ID:	P54652
Cytogenetics:	14q23.3
Domains:	HSP70
Protein Families:	Stem cell - Pluripotency
Protein Pathways:	Antigen processing and presentation, Endocytosis, MAPK signaling pathway, Spliceosome
Gene Summary:	Molecular chaperone implicated in a wide variety of cellular processes, including protection of the proteome from stress, folding and transport of newly synthesized polypeptides, activation of proteolysis of misfolded proteins and the formation and dissociation of protein complexes. Plays a pivotal role in the protein quality control system, ensuring the correct folding of proteins, the re-folding of misfolded proteins and controlling the targeting of proteins for subsequent degradation. This is achieved through cycles of ATP binding, ATP hydrolysis and ADP release, mediated by co-chaperones. The affinity for polypeptides is regulated by its nucleotide bound state. In the ATP-bound form, it has a low affinity for substrate proteins. However, upon hydrolysis of the ATP to ADP, it undergoes a conformational change that increases its affinity for substrate proteins. It goes through repeated cycles of ATP hydrolysis and nucleotide exchange, which permits cycles of substrate binding and release (PubMed:26865365). Plays a role in spermatogenesis. In association with SHCBP1L may participate in the maintenance of spindle integrity during meiosis in male germ cells (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RG200523