

## Product datasheet for **RG200538**

### PPP4C (NM\_002720) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	PPP4C (NM_002720) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	PPP4C
Synonyms:	PP-X; PP4; PP4C; PPH3; PPP4; PPX
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG200538 representing NM_002720 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCGGAGATCAGCGACCTGGACCGGCAGATCGAGCAGCTGCGTCGCTGCGAGCTCATCAAGGAGAGCG  
AAGTCAAGGCCCTGTGCGCTAAGGCCAGAGAGATCTTGGTAGAGGAGAGCAACGTGCAGAGGGTGGACTC  
GCCAGTCACAGTGTGCGGCGACATCCATGGACAATTCTATGACCTCAAAGAGCTGTTAGAGTAGGTGGC  
GACGTCCCTGAGACCAACTACCTCTTCATGGGGACTTTGTGGACCGTGGCTTCTATAGCGTCGAAACGT  
TCCTCCTGCTGCTGGCACTTAAGTTTCGCTATCCTGATCGCATCACACTGATCCGGGCAACCATGAGAG  
TCGCCAGATCACGAGGTCTATGGCTTCTACGATGAGTGCCTGCGCAAGTACGGCTCGGTGACTGTGTGG  
CGCTACTGCACTGAGATCTTTGACTACCTCAGCCTGTCAGCCATCATCGATGGCAAGATCTTCTGCGTGC  
ACGGGGGCTCTCCCCCTCCATCCAGACCCTGGATCAGATTCGGACAATCGACCGAAAGCAAGAGGTGCC  
TCATGATGGGCCCATGTGTGACCTCCTCTGGTCTGACCCAGAAGACACCACAGGCTGGGGCGTGAGCCCC  
CGAGGAGCCGGCTACCTATTTGGCAGTGACGTGGTGGCCAGTTCAACGCAGCCAATGACATTGACATGA  
TCTGCCGTGCCCACTACTGCTACCGCTGTGGAAATGTGGCAGCCATCTTGGAGCTGGACGAGCATCTCCAG  
AAAGATTCATCATCTTTGAGGCTGCTCCCCAAGAGACACGGGCATCCCCTCCAAGAAGCCCGTGGCCG  
ACTACTTCTG

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)


**Protein Sequence:** >RG200538 representing NM\_002720  
 Red=Cloning site Green=Tags(s)

MAEISDLDRQIEQLRRCELKSEVKALCAKAREILVEESNVQRVDSPTVCGDIHGQFYDLKELFRVGG  
 DVPETNYLFMGDFVDRGFYSVETFLLLLALKVRYPDRLTIRGNHESRQITQVYGFYDECLRKYGSVTW  
 RYCTEIFDYLSLSAIIIDGKIFCVHGGLSPSIQTLQDQIRTDKQEVPHDGPMDLLWSDPEDTTGWGVSP  
 RGAGYLFSGSDVVAQFNAANDIDMICRAHQQLVMEGYKWHFNETVLTVWSAPNYCYRCGNVAAILELDEHLQ  
 KDFIIFEAAPQETRGIPSKKPVADYFL

TRTRPLE - GFP Tag - V

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_002720

**ORF Size:** 921 bp


**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_002720.3</a>
<b>RefSeq Size:</b>	1429 bp
<b>RefSeq ORF:</b>	924 bp
<b>Locus ID:</b>	5531
<b>UniProt ID:</b>	<a href="#">P60510</a>
<b>Cytogenetics:</b>	16p11.2
<b>Domains:</b>	Metallophos, PP2Ac
<b>Protein Families:</b>	Druggable Genome, Phosphatase
<b>Gene Summary:</b>	<p>Protein phosphatase that is involved in many processes such as microtubule organization at centrosomes, maturation of spliceosomal snRNPs, apoptosis, DNA repair, tumor necrosis factor (TNF)-alpha signaling, activation of c-Jun N-terminal kinase MAPK8, regulation of histone acetylation, DNA damage checkpoint signaling, NF-kappa-B activation and cell migration. The PPP4C-PPP4R1 PP4 complex may play a role in dephosphorylation and regulation of HDAC3. The PPP4C-PPP4R2-PPP4R3A PP4 complex specifically dephosphorylates H2AFX phosphorylated on Ser-140 (gamma-H2AFX) generated during DNA replication and required for DNA double strand break repair. Dephosphorylates NDEL1 at CDK1 phosphorylation sites and negatively regulates CDK1 activity in interphase (By similarity). In response to DNA damage, catalyzes RPA2 dephosphorylation, an essential step for DNA repair since it allows the efficient RPA2-mediated recruitment of RAD51 to chromatin. [UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for RG200538