

Product datasheet for **RG200612**

EIF3B (NM_001037283) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	EIF3B (NM_001037283) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	EIF3B
Synonyms:	EIF3-ETA; EIF3-P110; EIF3-P116; EIF3S9; PRT1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RG200612 representing NM_001037283
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**GCGATCGCC**

ATGCAGGACGCGGAGAACGTGGCGGTGCCCGAGGCGGCCGAGGAGCGCGCCGAGCCCGCCAGCAGCAGC
 CGGCCCGAGCCGCGCCAGCCGAGGGGCTGCTGCGGCCCGCGGGCCCGGCGCTCCGGAGGCCGCGGG
 GACCGAGGCCCTCAGTGAGGAGGTGGGGATCGCGGAGGCCGCGGCCGAGCCCGAGGTGAGGACCGAGCCG
 GCGGCCGAGGCAGAGCGGCCCTCCGGCCCGTCCGAGTCCGCTCGCCCGCGGCCGCGCCGAGGAGCTGCCCG
 GGTGCGATGCTGAGCCCCCTGTCCGGCACAGGGCGAGGCCCCAGGAGAGCAGGCTCGGGACGAGCGCTC
 CGACAGCCGGGCCAGGCGGTGTCCGAGGACGCGGGAGGAAACGAGGGCAGAGCGGCCGAGGCCGAAACCC
 CGGGCGTGGAGAATGGCGACGCGGACGAGCCCTCCTTCAGCGACCCCGAGGACTTCGTGGACGACGTGA
 GCGAGGAAGAATTACTGGGAGATGTAACAAGATCGGCCCCAGGAAGCAGATGGAATCGATTCCGTTGAT
 TGTAGTGGACAATGCCCTCAGGTGGGACCCGACCGACTTGAGAACTCAAAAATGTCATCCACAAGATC
 TTTTCAAAGTTTGGGAAAATCACAAATGATTTTTATCCTGAAGAGGATGGGAAGCAAAAAGGGTATATTT
 TCCTGGAGTACGCGTCCCTGCCACGCTGTGGATGCTGTGAAGAACGCCGACGGCTACAAGCTTGACAA
 GCAGCACACATTCGGGTCAACCTCTTTACGGATTTTGACAAGTATATGACGATCAGTGACGAGTGGGAT
 ATTCCAGAGAAACAGCCTTTCAAAGACCTGGGGAACCTTACGTTACTGGCTTGAAGAGGCAGAATGCAGAG
 ATCAGTACAGTGTGATTTTTGAGAGTGGAGACCGCACTTCCATATTCTGGAATGACGTA AAAAGACCTGT
 CTAATTGAAGAAAGAGCGAGATGGACAGAGACGTATGTGCGTTGGTCTCCTAAGGGCACCTACCTGGCT
 ACCTTTTCATAAAGAGGCATTGCTCTATGGGGGGAGAGAAAATCAAGCAAATTCAGAGATTCAGCCACC
 AAGGGTTTCAGCTTATTGACTTTCACCTTGTGAAAGGTACCTGGTGACCTTTAGCCCTGATGGACAC
 GCAGGATGACCCTCAGGCCATAATCATCTGGGACATCCTTACGGGGCACAGAAGAGGGGTTTTCACTGT
 GAGAGCTCAGCCATTGGCCTATTTTTAAGTGGAGCCATGATGGCAAATCTTTGCCAGAATGACCTGG
 ATACGCTTAGCATCTATGAACTCCTTCTATGGGTCTTTTGGACAAGAAGAGTTTGAAGATCTCTGGGAT
 AAAAGACTTTTCTGGTCTCCTGGTGGTAACATAATCGCCTTCTGGTGCCTGAAGACAAAGATATTCCA
 GCCAGGGTAACCCTGATGCAGCTCCCTACCAGGCAAGAGATCCGAGTGAGGAACCTGTTCAATGTGGTGG
 ACTGCAAGCTCCATTGGCAGAAGAACGGAGACTACTTGTGTGAAAAGTAGATAGGACTCCGAAAGGCAC
 CCAGGGTGTGTACAAAATTTGAAATTTCCGAATGAGGGAGAAACAGGTACCTGTGGATGTGGTCCGAG
 ATGAAAGAAACCATCATAGCCTTTGCCCTGGGAACCAAATGGAAGTAAGTTTGTGTGCTGCACGGAGAGG
 CTCGCGGATATCTGTGCTTTCTACCAGCTCAAAAACAACGGGAAGATTGAACTCATCAAGATGTTTCA
 CAAGCAGCAGGCGAACACCATCTTCTGGAGCCCCAAGGACAGTTCGTGGTGTGGCGGGCTGAGGAGT
 ATGAACGGTGCCTTAGCGTTTGTGGACTTCCGACTGCACGGTGCATGAACATCGCAGAGCACTACATGG
 CTTCCGACGTGCAATGGGATCCTACTGGGCGCTACGTCGTACCTCTGTGCTCCTGGTGGAGCCATAAGGT
 GGACAACGCGTACTGGCTGTGGACTTCCAGGGACGCCTCCTGCAGAAGAACAACAAGGACCGCTTCTGC
 CAGCTGCTGTGGCGGCCCGCCCTCCACACTCCTGAGCCAGGAACAGATCAAGCAAATTA AAAAGGATC
 TGAAGAAATACTAAGATCTTTGAACAGAAGGATCGTTTGAGTCAGTCAAAGCCTCAAAGGAATTGGT
 GGAGAGAAGGCGCACCATGATGGAAGATTTCCGGAAGTACCGGAAAATGGCCAGGAGCTCTATATGGAG
 CAGAAAAACGAGCGCCTGGAGTTGCGAGGAGGGTGGACACTGACGAGCTGGACAGCAACGTGGACGACT
 GGAAGAGGAGACCATTGAGTTCTCGTACTGAAGAAATCATTCCCCTCGGGAATCAGGAG

AG**CGGACCG**ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG200612 representing NM_001037283
 Red=Cloning site Green=Tags(s)

```

MQDAENVAVPEAAEERAEPGQQPAAEPPPAEGLLRPAGPGAPEAAGTEASSEEVGIAEAGPEPEVRTEP
AAEAEAASGPSESPSPAAEELPGSHAEPVPAQGEAPGEQARDERSDSRAQAVSEDAGGNEGRAAEAEAP
RALENGDADEPSFSDPEDFVDDYSEEELLGDVLDKDRPQEAADGIDSVIVDNPVQVGPDRLEKLNVIHKI
FSKFGKITNDFYPEEDGKTKGYIFLEYASPAHAVDAVKNADGYKLDKQHTFRVNLFTDFDKYMTISDEWD
IPEKQPFKDLGNLRYWLEEAECRDQYSVIFESGDRTSIFWNDVKDPVSIERARWTETYVRWSPKGTyla
TFHQRGIALWGGEKFKQIQRFHQVQLIDFSPCERYLVTFSPMLDTQDDPQAIIIWDILTGHKRGRFHC
ESSAHPWIFKWSHDGKFFARMTLDTLSIYETPSMGLLDKSLKISGKDFSWSPGGNIIAFWVPEDKDIP
ARVTLMLQLPTRQEIRVRNLFNVVDCKLHWQKNGDYLCVKVDRTPKGTQGVVTFNFEIFRMREKQVPVDVVE
MKETIIAFWEPNGSKFAVLHGEAPRISVSFYHVKNNGKIELIKMFDKQANTIFWSPQGQFVVLAGLRS
MNGALAFVDTSDCTVMNIAEHYMASDVEWDPTGRYVVTSVSWWSHKVDNAYWLWTFQGRLLQKNNKDRFC
QLLWRPRPPTLLSQEQIKQIKKDLKKYSKIFEQKDRLSQSKASKELVERRRTMMEDFRKYRQMAQELYME
QKNERLELRGGVDTDELDSNVDDWEEETIEFFVTEEIIPLGNQE
  
```

SGPTRRRLE - GFP Tag - V

Restriction Sites: Sgfl-RsrII

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001037283.1](#), [NP_001032360.1](#)

RefSeq Size: 3084 bp

RefSeq ORF: 2445 bp


Locus ID: 8662

UniProt ID: [P55884](#)

Cytogenetics: 7p22.3

Gene Summary:

RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:9388245, PubMed:17581632, PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA_i and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:9388245, PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773). [UniProtKB/Swiss-Prot Function]

Product images:


Circular map for RG200612