

Product datasheet for **RG201248**

UAP56 (DDX39B) (NM_004640) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	UAP56 (DDX39B) (NM_004640) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	UAP56
Synonyms:	BAT1; D6S81E; UAP56
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG201248 representing NM_004640 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCAGAGAACGATGTGGACAATGAGCTCTTGGACTATGAAGATGATGAGGTGGAGACAGCAGCTGGGG
GAGATGGGGCTGAGGCCCTGCCAAGAAGGATGTCAAGGGCTCCTATGTCTCCATCCACAGCTCTGGCTT
TCGTGACTTCTGCTCAAGCCAGAGTTGCTCCGGGCCATTGTCGACTGTGGCTTTGAGCATCCGTCAGAA
GTCCAGCATGAGTGCATCCCTCAGGCCATTCTGGGAATGGATGTCCTGTGCCAGGCCAAGTCGGGCATGG
GAAAGACAGCAGTGTCTTGTCTTGGCCCACTGCAACAGCTGGAGCCAGTTACTGGCAGGTGTCTGTGCT
GGTGTGTGTCACTCGGGAGTTGGCTTTTCAGATCAGCAAGGAATATGAGCGCTTCTCTAAATACATG
CCCAATGTCAAGTTGCTGTTTTTTTTGGTGGTCTGTCTATCAAGAAGGATGAAGAGGTGCTGAAGAAGA
ACTGCCCGCATATCGTCGTGGGGACTCCAGGCCGTATCCTAGCCCTGGCTCGAAATAAGAGCCTCAACCT
CAAACACATTAACACTTTATTTTGGATGAATGTGATAAGATGCTTGAACAGCTCGACATGCGTCGGGAT
GTCCAGGAAATTTTTCGCATGACCCCCACGAGAAGCAGGTGATGATGTTCAAGTGTACCTTGAACAAAG
AGATCCGTCCAGTCTGCCCAAGTTCATGCAAGATCCAATGGAGATCTTCGTGGATGATGAGACGAAGTT
GACGCTGCATGGGTTGCAGCAGTACTACGTGAACTGAAGGACAACGAGAAGAACCGGAAGCTCTTTGAC
CTTCTGGATGTCCTTGAGTTCAACCAGGTGGTGTCTTTGTGAAGTCTGTGCAGCGGTGCATTGCCTTGG
CCAGCTACTAGTGGAGCAGAACTTCCCAGCCATTGCCATCCACCGTGGGATGCCCCAGGAGGAGAGGCT
TTCTCGGTATCAGCAGTTTAAAGATTTTCAACGACGAATCTTGTGGCTACCAACCTATTTGGCCGAGGC
ATGGACATCGAGCGGGTGAACATTGCTTTAATTATGACATGCCTGAGGATTCTGACACCTACCTGCATC
GGGTGGCCAGAGCAGGCCGTTTGGCACCAGGGCTTGGCTATCACATTTGTGTCCGATGAGAATGATGC
CAAGATCCTCAATGATGTGCAGGATCGCTTTGAGGTCAATATTAGTGAGCTGCCTGATGAGATAGACATC
TCCTCTACATTGAACAGACACGG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG201248 representing NM_004640
Red=Cloning site Green=Tags(s)

MAENDVDNELLDYEDDEVETAAGGDGAEAPAKKDVKGSYVSIHSSGFRDFLLKPELLRAIVDCGFEHPSE
 VQHECIPQAILGMDVLCQAKSGMGKTAVFVLATLQOLEPVTGQVSVLVMCHTRELAFQISKEYERFSKYM
 PNVKVAVFVFGGLSIKKDEEVLKKNCPHIVVGTGPRILALARNKSLNLKHIKHFILDECDKMLEQLDMRRD
 VQEIFRMTPEKQVMMFSATLSKEIRPVCRKFMQDPMEIFVDDETKLTLHGLQQYYVVKLDNEKNRKLFD
 LLDVLEFNQVVIFVKSQRCIALAQLLVEQNFPAAIAIHRGMPQEERLSRYQQFKDFQRRILVATNLFGRG
 MDIERNIAFNYPEDSDTYLHRVARAGRFGTKGLAITFVSDENDAKILNDVQDRFEVNI SELPDEIDI
 SSYIEQTR

TRTRPLE – GFP Tag – V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_004640

ORF Size: 1284 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_004640.4](#), [NP_004631.1](#)

RefSeq Size: 2133 bp

RefSeq ORF: 1287 bp

Locus ID: 7919

UniProt ID: [Q13838](#)


Cytogenetics: 6p21.33

Domains: DEAD, helicase_C

Protein Pathways: Spliceosome

Gene Summary: This gene encodes a member of the DEAD box family of RNA-dependent ATPases that mediate ATP hydrolysis during pre-mRNA splicing. The encoded protein is an essential splicing factor required for association of U2 small nuclear ribonucleoprotein with pre-mRNA, and it also plays an important role in mRNA export from the nucleus to the cytoplasm. This gene belongs to a cluster of genes localized in the vicinity of the genes encoding tumor necrosis factor alpha and tumor necrosis factor beta. These genes are all within the human major histocompatibility complex class III region. Mutations in this gene may be associated with rheumatoid arthritis. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on both chromosomes 6 and 11. Read-through transcription also occurs between this gene and the upstream ATP6V1G2 (ATPase, H⁺ transporting, lysosomal 13kDa, V1 subunit G2) gene. [provided by RefSeq, Feb 2011]

Product images:



Circular map for RG201248