

Product datasheet for RG201301

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

DDIT3 (NM 004083) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: DDIT3 (NM 004083) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: DDIT3

Synonyms: AltDDIT3; C/EBPzeta; CEBPZ; CHOP; CHOP-10; CHOP10; GADD153

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG201301 representing NM_004083

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

TGAATCTGCACCAAGCA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG201301 representing NM_004083

Red=Cloning site Green=Tags(s)

MAAESLPFSFGTLSSWELEAWYEDLQEVLSSDENGGTYVSPPGNEEEESKIFTTLDPASLAWLTEEEPEP AEVTSTSQSPHSPDSSQSSLAQEEEEEDQGRTRKRKQSGHSPARAGKQRMKEKEQENERKVAQLAEENER

LKQEIERLTREVEATRRALIDRMVNLHQA

TRTRPLE - GFP Tag - V


Restriction Sites: Sgfl-Mlul





Cloning Scheme:





ACCN: NM_004083

ORF Size: 507 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 004083.3</u>

RefSeq Size: 927 bp
RefSeq ORF: 510 bp
Locus ID: 1649
UniProt ID: P35638



Cytogenetics: 12q13.3 **Domains:** BRLZ

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: MAPK signaling pathway

Gene Summary: This gene encodes a member of the CCAAT/enhancer-binding protein (C/EBP) family of

> transcription factors. The protein functions as a dominant-negative inhibitor by forming heterodimers with other C/EBP members, such as C/EBP and LAP (liver activator protein), and

preventing their DNA binding activity. The protein is implicated in adipogenesis and


erythropoiesis, is activated by endoplasmic reticulum stress, and promotes apoptosis. Fusion

of this gene and FUS on chromosome 16 or EWSR1 on chromosome 22 induced by

translocation generates chimeric proteins in myxoid liposarcomas or Ewing sarcoma. Multiple alternatively spliced transcript variants encoding two isoforms with different length have

been identified. [provided by RefSeq, Aug 2010]

Product images:



Circular map for RG201301