

## Product datasheet for **RG201314**

### **Bcl x (BCL2L1) (NM\_138578) Human Tagged ORF Clone**

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Bcl x (BCL2L1) (NM_138578) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Bcl x
Synonyms:	Bcl-X; BCL-XL/S; BCL2L; BCLX; PPP1R52
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG201314 representing NM_138578 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGTCTCAGAGCAACCGGGAGCTGGTGGTTGACTTTCTCTCCTACAAGCTTTCCAGAAAGGATACAGCT  
GGAGTCAGTTTAGTGATGTGGAAGAGAACAGGACTGAGGCCCCAGAAGGGACTGAATCGGAGATGGAGAC  
CCCCAGTGCCATCAATGGCAACCCATCCTGGCACCTGGCAGACAGCCCCGGTGAATGGAGCCACTGGC  
CACAGCAGCAGTTTGGATGCCCGGGAGGTGATCCCATGGCAGCAGTAAAGCAAGCGCTGAGGGAGGCAG  
GCGACGAGTTTGAAGTACCGGTACCGCGGGCATTGAGTACCTGACATCCAGCTCCACATCACCCAGG  
GACAGCATATCAGAGCTTTGAACAGGTAGTGAATGAACTTCCGGGATGGGGTAAACTGGGGTCCGATT  
GTGGCCTTTTTCTCCTTCGGCGGGCACTGTGCGTGAAAAGCGTAGACAAGGAGATGCAGGTATTGGTGA  
GTCGGATCGCAGCTTGGATGGCCACTTACCTGAATGACCACCTAGAGCCTTGGATCCAGGAGAACGGCGG  
CTGGGATACTTTTGTGGAAGTCTATGGGAACAATGCAGCAGCCGAGAGCCGAAAGGGCCAGGAACGCTTC  
AACCGTGGTTCTGACGGGCATGACTGTGGCCGGCGTGGTTCTGCTGGGCTCACTTTCAGTCGGAAA

**ACGCGT**ACGCGGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG201314 representing NM\_138578  
Red=Cloning site Green=Tags(s)

MSQSNRELVVDFLSYKLSQKGYWSQFSDVEENRTEAPEGTESEMETPSAINGNPSWHLADSPAVNGATG  
 HSSSLDAREVIPMAAVKQALREAGDEFELRYRRAFSDLTSQLHITPGTAYQSFEQVYNELFRDGVNWGRI  
 VAFFSFGGALCVESVDKEMQVLVSRIAAMATYLNHLEPWIQENGGWDTFVELYGNAAAESRKGQERF  
 NRWFLTGMTVAGVVLLGSLFSRK

TRTRPLE - GFP Tag - V

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_138578

**ORF Size:** 699 bp


**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_138578.3</a>
<b>RefSeq Size:</b>	2575 bp
<b>RefSeq ORF:</b>	702 bp
<b>Locus ID:</b>	598
<b>UniProt ID:</b>	<a href="#">Q07817</a>
<b>Cytogenetics:</b>	20q11.21
<b>Domains:</b>	Bcl-2, BH4
<b>Protein Families:</b>	Druggable Genome, ES Cell Differentiation/IPS, Transmembrane
<b>Protein Pathways:</b>	Amyotrophic lateral sclerosis (ALS), Apoptosis, Chronic myeloid leukemia, Jak-STAT signaling pathway, Pancreatic cancer, Pathways in cancer, Small cell lung cancer
<b>Gene Summary:</b>	<p>The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform acts as an apoptotic inhibitor and the shorter isoform acts as an apoptotic activator. [provided by RefSeq, Dec 2015]</p>

Product images:



Circular map for RG201314