

Product datasheet for **RG201822**

DUSP11 (NM_003584) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	DUSP11 (NM_003584) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	DUSP11
Synonyms:	PIR1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG201822 representing NM_003584 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAGCCAGTGGCATCATCCCCGAGTGGCTGGGGCCGGAGACGCGACTTTTCAGGACGCTCCTCAGCCA
AGAAGAAGGGCGGAAACCACATCCCCGAAAGGTGGAAAGACTATCTCCAGTTGGACAGCGGATGCCTGG
GACTCGTTTCATTGCTTTCAAAGTTCCTTTGCAAAGAGTTTTGAAAAGAACTTGCTCCAGAAGAATGC
TTTTCCCTTTGGATCTTTTAAACAAAATCCGAGAACAAAATGAAGAAGTGGACTGATTATTGATTTAA
CATATACTCAACGCTATTATAAACCAGAGGATTTGCCAGAACTGTTCTTACTTAAAAATTTTACAGT
TGGACATCAAGTGCCTGATGATGAGACTATTTTTAAATTCAAACACGCTGTTAATGGGTTTTTGAAGAA
AATAAAGATAATGATAAACTTATTGGTGTCCACTGTACCCATGGTTTAAACAGGACTGGCTACCTCATT
GCATATATTTGATTGATGTAGAAGGCGTGAGGCCAGATGATGCAATTGAATTATTCAATAGGTGCCGGGG
ACATTGCTTAGAAAGACAAAACACTACATTGAAGACCTTCAGAAATGGTCCTATCAGAAAGAATTGGAATTCC
AGTGTACCCAGGTCAAGTATTTGAAGACTCAGCACATCTCATGCAACCCAGTCCACAATAAGCCTGTTA
ACAAGGACCTAGGTATAATCTACATCAGATCCAGGGTCACTCAGCTCCTCGACATTTCCACACCCAGAC
CCAAAGTTTGAACAATCAGTCAGAAAAATTTTCAGAGAATCCACATGTTTACCAGAGACACCATCTCCCT
CCTCCTGGTCCCCCTGGAGAGGACTATTCACACAGGAGTATTCTTGAATGTGAAGCCCAATGCCAGTC
GGGCAGCCAGGATAGAAGAAGGTGGTATCCTTATAATTACTCCAGACTCTCCTATCCAGCCTGTTGGGA
ATGGACCCAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG201822 representing NM_003584
 Red=Cloning site Green=Tags(s)

MSQWHHRSGWGRRRDFSGRSSAKKKGGNHIPERWKDYLPVQQRMPGTRFIAFKVPLQKSFEKLAPEEC
 FSPLDLFNKIREQNEELGLIIDLTYTQRYYPEDLPETVPYLKIFTVGHQVPDDETFKFKHAVNGFLKE
 NKDNDKLGIVHCTHGLNRTGYLICIYLIDVEGVRPDDAIELFNRCRGHCLERQNYIEDLQNGPIRKNWNS
 SVPRSSDFEDSAHLMQPVHNKPKVQGPVYLNHQIQGHSAPRHFHTQTQSLQQSVRKFSENPHVYQRHHLPP
 PGGPPGEDYSHRRYSWNVKNASRAAQRDRRRWYPYNYSRLSYPACWEWTQ

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_003584

ORF Size: 990 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_003584.1 , NP_003575.1
RefSeq Size:	1593 bp
RefSeq ORF:	1134 bp
Locus ID:	8446
UniProt ID:	O75319
Cytogenetics:	2p13.1
Domains:	DSPc
Protein Families:	Druggable Genome, Phosphatase
Gene Summary:	<p>The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which is associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product is localized to the nucleus and binds directly to RNA and splicing factors, and thus it is suggested to participate in nuclear mRNA metabolism. [provided by RefSeq, Sep 2008]</p>

Product images:



Circular map for RG201822