

## Product datasheet for **RG201950**

### ELOC (NM\_005648) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** ELOC (NM\_005648) Human Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** ELOC  
**Synonyms:** SIII; TCEB1  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >RG201950 representing NM\_005648  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGGATGGAGAGGAGAAAACCTATGGTGGCTGTGAAGGACCTGATGCCATGTATGTCAAATTGATATCAT  
 CTGATGGCCATGAATTTATTGTAAAAAGAGAACATGCATTAACATCAGGCACGATAAAAGCCATGTTGAG  
 TGGCCAGGTCAAGTTGCTGAGAACGAAACCAATGAGGTCAATTTAGAGAGATACCTTCACATGTGCTA  
 TCGAAAGTATGCATGTATTTACGTACAAGTTCGCTACACTAACAGCTCCACCGAGATTCCTGAATTCC  
 CAATTGCACCTGAAATTGCACTGGAAGTCTGATGGCTGCGAAGTCTTAGATTGT

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >RG201950 representing NM\_005648  
 Red=Cloning site Green=Tags(s)  
 MDGEEKTYGGCEGPDAMYVKLISSDGHEFIVKREHALTSGTIKAMLSGPGQFAENETNEVNFREIPSHVL  
 SKVCMYFTYKVRVTNSSTEIPEFPPIAPEIALELLMAANFLDC

**TRTRPLE** - GFP Tag - V

**Restriction Sites:** Sgfl-MluI



[View online »](#)

**Cloning Scheme:**


**ACCN:** NM\_005648

**ORF Size:** 336 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
  2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
  3. Close the tube and incubate for 10 minutes at room temperature.
  4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
  5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.


**RefSeq:** [NM\\_005648.4](#)

**RefSeq Size:** 668 bp

**RefSeq ORF:** 339 bp

**Locus ID:** 6921  
**UniProt ID:** [Q15369](#)  
**Cytogenetics:** 8q21.11  
**Domains:** Skp1  
**Protein Families:** Druggable Genome, Transcription Factors  
**Protein Pathways:** Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysis  
**Gene Summary:** This gene encodes the protein elongin C, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been identified. [provided by RefSeq, Mar 2011]

**Product images:**



Circular map for RG201950