

Product datasheet for **RG203771**

DAP12 (TYROBP) (NM_198125) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: DAP12 (TYROBP) (NM_198125) Human Tagged ORF Clone
Tag: TurboGFP
Symbol: DAP12
Synonyms: DAP12; KARAP; PLOSL; PLOSL1
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >RG203771 representing NM_198125
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGGGGACTTGAACCTGCAGCAGGCTCCTGCTCCTGCCTCCTGCTGGCTGTAAGTGGTCTCCGTC
CTGTCCAGGCCAGGCCAGAGCGATTGCAGTTGCTCTACGGTGAGCCCGGGCGTCTGGCAGGGATCGT
GATGGGAGACCTGGTCTGACAGTGCTCATTGCCCTGGCCGTGTACTTCTGGCCGGCTGGTCCCTCGG
GGCGGAGGGCTGCGGAGGCAGCGACCCGAAACAGCGTATCACTGAGACCGAGTCGCCTTATCAGGAGC
TCCAGGGTCAGAGGTCGGATGTCTACAGCGACCTCAACACACAGAGGCCGTATTACAAA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA


Protein Sequence: >RG203771 representing NM_198125
Red=Cloning site Green=Tags(s)
MGGLEPCSRLLLLPLLLAVSGLRPVQAQAQSDCSCSTVSPGVLAGIVMGDLVLTVLIALAVYFLGRLLVPR
GRGAAEAATRKQRITETESPYQELQQRSDVYSDLNTQRPYYK

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-MluI



[View online »](#)

Cloning Scheme:


ACCN: NM_198125

ORF Size: 339 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_198125.1](#), [NP_937758.1](#)


RefSeq Size: 580 bp

RefSeq ORF: 339 bp

Locus ID: 7305
UniProt ID: [O43914](#)
Cytogenetics: 19q13.12
Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Natural killer cell mediated cytotoxicity
Gene Summary:

This gene encodes a transmembrane signaling polypeptide which contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. The encoded protein may associate with the killer-cell inhibitory receptor (KIR) family of membrane glycoproteins and may act as an activating signal transduction element. This protein may bind zeta-chain (TCR) associated protein kinase 70kDa (ZAP-70) and spleen tyrosine kinase (SYK) and play a role in signal transduction, bone modeling, brain myelination, and inflammation. Mutations within this gene have been associated with polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy (PLOSL), also known as Nasu-Hakola disease. Its putative receptor, triggering receptor expressed on myeloid cells 2 (TREM2), also causes PLOSL. Multiple alternative transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Mar 2010]

Product images:



Circular map for RG203771