

Product datasheet for **RG204159**

P2Y2 (P2RY2) (NM_002564) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: P2Y2 (P2RY2) (NM_002564) Human Tagged ORF Clone
Tag: TurboGFP
Symbol: P2RY2
Synonyms: HP2U; P2RU1; P2U; P2U1; P2UR; P2Y2; P2Y2R
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >RG204159 representing NM_002564
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGCAGCAGACCTGGGCCCTGGAATGACACCATCAATGGCACCTGGGATGGGGATGAGCTGGGCTACA
 GGTGCCGCTTCAACGAGGACTTCAAGTACGTGCTGCTGCCTGTGTCTACGGCGTGGTGTGCGTGTGG
 GCTGTGTCTGAACGCCGTGGCGCTCTACATCTTCTGTGCCGCTCAAGACCTGGAATGCGTCCACCACA
 TATATGTTCCACCTGGCTGTGTCTGATGACTGTATGCGGCTCCCTGCCGCTGCTGGTCTATTACTACG
 CCCGGCGGACCACTGGCCCTTACGACGGTCTGCAAGCTGGTGGCTTCTCTTCTACACCAACCT
 TTAAGTGCATCCTTCTCCTCAGTGCATCAGCGTGCACCGGTGCTGGGCGTCTTACGACCTCTGCGC
 TCCTGCGCTGGGGCCGGCCCGCTACGCTCGCCGGTGGCCGGGGCCGTGTGGTGTGGTGTGGCT
 GCCAGGCCCGTGTCTACTTTGTACCAACAGCGCGCGGGGGCCGTAACCTGCCACGACACCTC
 GGCACCCGAGCTTTCAGCCGCTTCGTGGCCTACAGCTCAGTCATGCTGGCCTGCTTTCGCGGTGCC
 TTTGCCGTCATCCTTGTCTGTTACGTGCTATGGCTCGGCGACTGCTAAAGCCAGCTACGGGACCTCGG
 GCGGCTGCCTAGGGCAAGCGCAAGTCCGTGCGCACCATCGCGTGGTGTGCTGGTGTCTTCGCCCTCG
 CTTCTGCCATTCACGTACCCGACCCCTCTACTACTCCTTCCGTTTCGTTGGACCTCAGCTGCCACCC
 CTCACGCCATCAACATGGCTACAAGTTACCCGGCCGCTGGCCAGTGTAAAGTTGCCTTGACCCCG
 TGCTCTACTTCTGGCTGGGACAGCCTCGTACGCTTTGCCGAGATGCCAAGCCACCCACTGGCCCCAG
 CCCTGCCACCCCGCTCGCCGAGGCTGGGCTGCGCAGATCCGACAGAAGTGCATGCAGAGGATAGAA
 GATGTGTTGGCAGCAGTGGGACTCTAGGCGGACAGAGTCCACGCCGGTGGTAGCGAGAACAATAAGG
 ACATTCGGCTG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG204159 representing NM_002564
 Red=Cloning site Green=Tags(s)

MAADLGPWNDDINGTWDGDELGYRCRFNEDFKYVLLPVSYGVVVCLGLCLNAVALYIFLCRLKTNASTT
 YMFHLAVSDALYAASPLLVYYYARGDHPFSTVLCKLVRFLFYTNLYCSILFLTCISVHRCLGVLRLR
 SLRWGRARYARRVAGAVWLVLACQAPVLYFVTT SARGGRVTCHDTSAPELFSRFVAYSSVMLGLLFAVP
 FAVILVCYVLMARRLLKPAYGTSGGLPRAKRKSVRTIAVVLAVFALCFLPFHVTRTLYYFSRSLDLSCHT
 LNAINMAYKVTRPLASANSCLDPVLYFLAGQSLVRFARDAKPPTGSPATPARRRLGLRRSDRTDMQRIE
 DVLGSSSEDSRRTTESTPAGESNTKDIRL


TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_002564

ORF Size:	1131 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_002564.2 , NP_002555.2
RefSeq Size:	2547 bp
RefSeq ORF:	1134 bp
Locus ID:	5029
UniProt ID:	P41231
Cytogenetics:	11q13.4
Domains:	7tm_1
Protein Families:	Druggable Genome, GPCR, Transmembrane
Protein Pathways:	Neuroactive ligand-receptor interaction
Gene Summary:	The product of this gene belongs to the family of P2 receptors, which is activated by extracellular nucleotides and subdivided into P2X ligand-gated ion channels and P2Y G-protein coupled receptors. This family has several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor, found on many cell types, is activated by ATP and UTP and is reported to be overexpressed on some cancer cell types. It is involved in many cellular functions, such as proliferation, apoptosis and inflammation. Three transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq, Mar 2013]