

## Product datasheet for **RG204685**

### ATP6V1D (NM\_015994) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	ATP6V1D (NM_015994) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	ATP6V1D
Synonyms:	ATP6M; VATD; VMA8
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG204685 representing NM_015994 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGTCGGGCAAAGACCGAATTGAAATCTTCCCTCGGAATGGCACAGACCATCATGAAGGCTCGTTTAA  
AGGGAGCACAGACAGGTCGAAACCTCCTGAAGAAAAATCTGATGCCTTAACTCTTCGATTTGACAGAT  
CCTAAAGAAGATAATAGAGACTAAAATGTTGATGGCGAAGTGATGAGAGAAGCTGCCTTTTCACTAGCT  
GAAGCCATGTTTACAGCAGGTGACTTCAGCACTACAGTTATCCAAAATGTCAATAAAGCGCAAGTGAAGA  
TTCGAGCGAAGAAAGATAATGTAGCAGGTGTTACTTTGCCAGTATTTGAACATTACCATGAAGGAAGTGA  
CAGTTATGAAGTACTGTTTAGCCAGAGGTGGGGAACAGTTGGCTAAATTAAGAGGAATTATGCCAAA  
GCAGTGGAACTACTGGTGGAACTAGCTTCTCTGCAGACTTCTTTTGTACTTTGGATGAAGCTATTAAGA  
TAACCAACAGGCGTGAAATGCCATTGAACATGTCATCATTCCCGGATTGAACGTAAGTCTTGTCTATAT  
CATCACAGAGCTGGATGAGAGAGAGCGAGAAGAGTTCTATAGGTTAAAGAAAATACAAGAGAAGAAAAAG  
ATTCTAAAGAAAAATCTGAGAAGGACTTGGAGCAAAGGAGAGCAGCTGGAGAGGTGTTGGAGCCTGCTA  
ATCTTCTGGCTGAAGAGAAGGACGAGGATCTTCTATTTGAA

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG204685 representing NM\_015994  
 Red=Cloning site Green=Tags(s)

```
MSGKDRIEIFPSRMAQTIMKARLKGAQTGRNLLKKKSDALTLRFRQILKKI IETKMLMGEVMREAAFSLA
EAMFTAGDFSTTVIQNVNKAQVKIRAKKDNVAGVTLPVFEHYHEGTD SYELTGLARGGEQLAKLKRNYAK
AVELLVELASLQTSFVTLDEAIKITNRRVNAIEHVIIIPRIERTLAYIITELDEREREEFYRLKKIQEKKK
ILKEKSEKDLQRRRAAGEVLEPANLLAEKDEDLLFE
```

TRTRPLE - GFP Tag - V

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_015994

**ORF Size:** 741 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_015994.2](#), [NP\\_057078.1](#)

**RefSeq Size:** 1675 bp

**RefSeq ORF:** 744 bp

**Locus ID:** 51382

**UniProt ID:** [Q9Y5K8](#)


**Cytogenetics:** 14q23.3

**Domains:** ATP-synt\_D

**Protein Pathways:** Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection

**Gene Summary:** This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c", and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This gene encodes the V1 domain D subunit protein. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RG204685