

Product datasheet for RG204821

C11orf10 (TMEM258) (NM_014206) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Tag: TurboGFP

Symbol: Cllorf10

Synonyms: Cllorf10; Kud; Kuduk

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide Sequence: >RG204821 representing NM_014206

Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGAGCTCGAGGCCATGAGCAGATATACCAGCCCAGTGAACCCAGCTGTCTTCCCCCATCTGACCGTGG
TGCTTTTTGGCCATTGGCATGTTCTTCACCGCCTGGTTCTTCGTTTACGAGGTCACCTCTACCAAGTACAC
TCGTGATATCTATAAAGAGCTCCTCATCTCCTTAGTGGCCTCACCTCTTCATGGGCTTTGGAGTCCTCTTC

CTGCTGCTCTGGGTTGGCATCTACGTG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG204821 representing NM_014206

Red=Cloning site Green=Tags(s)

MELEAMSRYTSPVNPAVFPHLTVVLLAIGMFFTAWFFVYEVTSTKYTRDIYKELLISLVASLFMGFGVLF

LLLWVGIYV

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn



Cloning Scheme:





ACCN: NM_014206

ORF Size: 237 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

C11orf10 (TMEM258) (NM_014206) Human Tagged ORF Clone | RG204821

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um

filter is required.

RefSeq: <u>NM_014206.4</u>

RefSeq Size: 418 bp

RefSeq ORF: 240 bp

Locus ID: 746

UniProt ID: P61165

Cytogenetics: 11q12.2

Protein Families: Transmembrane

Gene Summary: Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a

defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-

pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent

polypeptide chains, the first step in protein N-glycosylation. N-glycosylation occurs


cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum

(ER). All subunits are required for a maximal enzyme activity (PubMed:26472760,

PubMed:27974209). Involved in ER homeostasis in the colonic epithelium (By similarity).

[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RG204821