

### Product datasheet for RG204891

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Deoxyguanosine kinase (DGUOK) (NM\_080916) Human Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

Product Name: Deoxyguanosine kinase (DGUOK) (NM\_080916) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: Deoxyguanosine kinase

Synonyms: dGK; MTDPS3; NCPH; PEOB4

Mammalian Cell

Selection:

Neomycin

**Vector:** pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG204891 representing NM\_080916

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA





Protein Sequence: >RG204891 representing NM\_080916

Red=Cloning site Green=Tags(s)


MAAGRLFLSRLRAPFSSMAKSPLEGVSSSRGLHAGRGPRRLSIEGNIAVGKSTFVKLLTKTYPEWHVATE PVATWQNIQAAGTQKACTAQSLGNLLDMMYREPARWSYTFQTFSFLSRLKVQLEPFPEKLLQARKPVQIF ERSVYSDRYIFAKNLFENGSLSDIEWHIYQDWHSFLLWEFASRITLHGFIYLQASPQVCLKRLYQRAREE EKGIELAYLEQLHGQHEAWLIHKTTKLHFEALMNIPVLVLDVNDDFSEEVTKQEDLMREVNTFVKNL


TRTRPLE - GFP Tag - V

**Restriction Sites:** 

Sgfl-Mlul

**Cloning Scheme:** 





**ACCN:** NM\_080916

ORF Size: 831 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



#### **Reconstitution Method:**

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>NM 080916.3</u>

 RefSeq Size:
 1144 bp

 RefSeq ORF:
 834 bp

 Locus ID:
 1716

 UniProt ID:
 Q16854

 Cytogenetics:
 2p13.1

**Protein Families:** Druggable Genome


**Protein Pathways:** Metabolic pathways, Purine metabolism

**Gene Summary:** In mammalian cells, the phosphorylation of purine deoxyribonucleosides is mediated

predominantly by two deoxyribonucleoside kinases, cytosolic deoxycytidine kinase and mitochondrial deoxyguanosine kinase. The protein encoded by this gene is responsible for phosphorylation of purine deoxyribonucleosides in the mitochondrial matrix. In addition, this protein phosphorylates several purine deoxyribonucleoside analogs used in the treatment of lymphoproliferative disorders, and this phosphorylation is critical for the effectiveness of the analogs. Alternative splice variants encoding different protein isoforms have been described

for this gene. [provided by RefSeq, Jul 2008]

# **Product images:**



Circular map for RG204891