

Product datasheet for **RG204960**

RPLP0 (NM_001002) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	RPLP0 (NM_001002) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	RPLP0
Synonyms:	L10E; LP0; P0; PRLP0; RPP0
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG204960 representing NM_001002 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCCAGGGAAGACAGGGCGACCTGGAAGTCCAACACTTCTTAAGATCATCCAACATTGGATGATT
ATCCGAAATGTTTCATTGTGGGAGCAGACAATGTGGGCTCCAAGCAGATGCAGCAGATCCGCATGTCCT
TCGCGGGAAGGCTGTGGTGTGATGGGCAAGAACCATTGATGCGCAAGGCCATCCGAGGGCACCTGGAA
AACAAACCAGCTCTGGAGAACTGCTGCCTCATATCCGGGGAAATGTGGGCTTTGTGTTACCAAGGAGG
ACCTCACTGAGATCAGGGACATGTTGCTGGCCAATAAGGTGCCAGCTGCTGCCCGTCTGGTGCCATTGC
CCCATGTGAAGTCACTGTGCCAGCCAGAACACTGGTCTCGGGCCGAGAAGACCTCCTTTTTCCAGGCT
TTAGGTATCACCACTAAAATCTCCAGGGGCACCATTGAAATCCTGAGTGTGTGCAGCTGATCAAGACTG
GAGACAAAGTGGGAGCCAGCGAAGCCACGCTGCTGAACATGCTCAACATCTCCCCCTTCTCTTTGGGCT
GGTCATCCAGCAGGTGTTTCGACAATGGCAGCATCTACAACCTGAAGTGCTTGATATCACAGAGGAACT
CTGCATTCTCGCTTCTGGAGGGTGTCCGCAATGTTGCCAGTGTCTGTCTGCAGATTGGCTACCCAACTG
TTGCATCAGTACCCATTCTATCATCAACGGGTACAAACGAGTCTGGCCTTGTCTGTGGAGACGGATTA
CACCTTCCCACTTGCTGAAAAGGTCAAGGCCTTCTGGCTGATCCATCTGCCTTTGTGGCTGCTGCCCT
GTGGCTGCTGCCACCACAGCTGCTCCTGCTGCTGCTGCAGCCCAAGGTTGAAGCCAAGGAAGAGT
CGGAGGAGTCGGACGAGGATATGGGATTTGGTCTCTTTGAC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG204960 representing NM_001002
 Red=Cloning site Green=Tags(s)

MPREDRATWKSNYFLKIIQLDDYPKCFIVGADNVGSKQMQQIRMSLRGKAVVLMGKNTMMRKAIRGHLE
 NNPALAKLLPHIRGNVGFVFTKEDLTEIRDMLLANKVPAAARAGAIAPCEVTVP AQNTGLGPEKTSFFQA
 LGITTKISRGTIEILSDVQLIKTGDKVGASEATLLNMLNISPFSFGLVIQQVFDNGSIYNPEVLDITEET
 LHSRFLEGVRNVASVCLQIGYPTVASVPHSIINGYKRVLALSVETDYTFPLAEKVKAFADPSAFVAAAP
 VAAATTAAPAAAAAPAKVEAKEESESEDEDMGFLFD

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: NM_001002

ORF Size: 951 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001002.4
RefSeq Size:	1229 bp
RefSeq ORF:	954 bp
Locus ID:	6175
UniProt ID:	P05388
Cytogenetics:	12q24.23
Domains:	Ribosomal_L10, 60s_ribosomal
Protein Pathways:	Ribosome
Gene Summary:	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein, which is the functional equivalent of the E. coli L10 ribosomal protein, belongs to the L10P family of ribosomal proteins. It is a neutral phosphoprotein with a C-terminal end that is nearly identical to the C-terminal ends of the acidic ribosomal phosphoproteins P1 and P2. The P0 protein can interact with P1 and P2 to form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located in the cytoplasm. Transcript variants derived from alternative splicing exist; they encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]