

## Product datasheet for **RG205092**

### **RACK1 (NM\_006098) Human Tagged ORF Clone**

#### Product data:

Product Type:	Expression Plasmids
Product Name:	RACK1 (NM_006098) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	RACK1
Synonyms:	Gnb2-rs1; GNB2L1; H12.3; HLC-7; PIG21
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG205092 representing NM_006098 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGACTGAGCAGATGACCCTTCGTGGCACCCCTCAAGGGCCACAACGGCTGGGTAACCCAGATCGCTACTA  
CCCCGAGTCCCGGACATGATCCTCTCCGCTCTCGAGATAAGACCATCATCATGTGGAACTGACCAG  
GGATGAGACCAACTATGGAATCCACAGCGTCTCTCGGGGTCCTCCACTTTGTTAGTGATGTGGTT  
ATCTCTCAGATGGCCAGTTTCCCTCTCAGGCTCCTGGGATGGAACCTGCGCCTCTGGGATCTCACAA  
CGGGCACCACCAGAGGCGATTTGTGGCCATACCAAGGATGTGCTGAGTGTGGCCTTCTCCTCTGACAA  
CCGGCAGATTGTCTCTGGATCTCGAGATAAAACCATCAAGCTATGGAATACCCCTGGGTGTGTGCAAATAC  
ACTGTCCAGGATGAGAGCCACTCAGAGTGGGTGTCTTGTGTCCGCTTCTCGCCAAACAGCAGCAACCCCTA  
TCATCGTCTCCTGTGGCTGGGACAAGCTGGTCAAGGTATGGAACCTGGCTAACTGCAAGCTGAAGACCAA  
CCACATTGGCCACACAGGCTATCTGAACACGGTACTGTCTCTCCAGATGGATCCCTCTGTGCTTCTGGA  
GGCAAGGATGGCCAGGCCATGTTATGGGATCTCAACGAAGGCAAACACCTTTACACGCTAGATGGTGGG  
ACATCATCAACGCCCTGTGCTTCAGCCCTAACCCTACTGGCTGTGTGCTGCCACAGGCCCCAGCATCAA  
GATCTGGGATTTAGAGGGAAAGATCATTGTAGATGAACTGAAGCAAGAAGTTATCAGTACCAGCAGCAAG  
GCAGAACCACCCAGTGCACCTCCCTGGCTGGTCTGCTGATGGCCAGACTCTGTTTGTGGCTACACGG  
ACAACCTGGTGGAGTGTGGCAGGTGACCATTGGCACACGC

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)


**Protein Sequence:** >RG205092 representing NM\_006098  
 Red=Cloning site Green=Tags(s)

MTEQMTLRGTLKGHNGWVTQIATTPQFPDMILSASRDKTIIMWKLTRDETNYGIPQRALRGHSHFVSDVV  
 ISSDGFALSGSWDGLRLWDLTTGTTTRRFVGHTKDVLVAVFSSDNRQIVSGSRDKTIKLWNTLVGCKY  
 TVQDESHSEWVSCVRFSPNSSNP IIVSCGWDKLKVVWNLANCKLKTNHIGHTGYLNTVTVSPDGLSCASG  
 GKDGQAMLWDLNEGKHL YTLDDGGDIINALCFSPNRYWLC AATGPSIKIWDLEGKIIIVDELKQEVISTSSK  
 AEPPQCTSLAWSADGQTLFAGYTDNLVVRVWQVTIGTR

TRTRPLE - GFP Tag - V

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_006098

**ORF Size:** 951 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_006098.5](#)

**RefSeq Size:** 1075 bp

**RefSeq ORF:** 954 bp

**Locus ID:** 10399

**UniProt ID:** [P63244](#)


**Cytogenetics:** 5q35.3

**Domains:** WD40

**Protein Families:** Druggable Genome

**Gene Summary:** Scaffolding protein involved in the recruitment, assembly and/or regulation of a variety of signaling molecules. Interacts with a wide variety of proteins and plays a role in many cellular processes. Component of the 40S ribosomal subunit involved in translational repression (PubMed:23636399). Involved in the initiation of the ribosome quality control (RQC), a pathway that takes place when a ribosome has stalled during translation, by promoting ubiquitination of a subset of 40S ribosomal subunits (PubMed:28132843). Binds to and stabilizes activated protein kinase C (PKC), increasing PKC-mediated phosphorylation. May recruit activated PKC to the ribosome, leading to phosphorylation of EIF6. Inhibits the activity of SRC kinases including SRC, LCK and YES1. Inhibits cell growth by prolonging the G0/G1 phase of the cell cycle. Enhances phosphorylation of BMAL1 by PRKCA and inhibits transcriptional activity of the BMAL1-CLOCK heterodimer. Facilitates ligand-independent nuclear translocation of AR following PKC activation, represses AR transactivation activity and is required for phosphorylation of AR by SRC. Modulates IGF1R-dependent integrin signaling and promotes cell spreading and contact with the extracellular matrix. Involved in PKC-dependent translocation of ADAM12 to the cell membrane. Promotes the ubiquitination and proteasome-mediated degradation of proteins such as CLEC1B and HIF1A. Required for VANGL2 membrane localization, inhibits Wnt signaling, and regulates cellular polarization and oriented cell division during gastrulation. Required for PTK2/FAK1 phosphorylation and dephosphorylation. Regulates internalization of the muscarinic receptor CHRM2. Promotes apoptosis by increasing oligomerization of BAX and disrupting the interaction of BAX with the anti-apoptotic factor BCL2L. Inhibits TRPM6 channel activity. Regulates cell surface expression of some GPCRs such as TBXA2R. Plays a role in regulation of FLT1-mediated cell migration. Involved in the transport of ABCB4 from the Golgi to the apical bile canalicular membrane (PubMed:19674157). Promotes migration of breast carcinoma cells by binding to and activating RHOA (PubMed:20499158).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RG205092