

Product datasheet for **RG207172**

Protocadherin beta 12 (PCDHB12) (NM_018932) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Tag:	TurboGFP
Symbol:	Protocadherin beta 12
Synonyms:	PCDH-BETA12
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



ORF Nucleotide Sequence: >RG207172 representing NM_018932
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGC**

ATGGAACCGAGGGGCGAGCACTCTGCAGATAAGGCAAGTCTGCTTTCTTTGTTTGTGGGAATGT
CTCAGGCGGGCTCTGAACTGGGAACTTTTGGTGATGGAGGAATTGCAGAGCGGGAGCTTTGTAGGAAA
TTTGGCAAAGACCTGGGACTCGAGGTGAGTGAGCTGTCTTCGCGGGGGCTCGGGTGGTTTCTAATGAT
AACAAGAGTGTTTGCAGCTGGACACAACTGGGATTTGCTCCTGAGAGAAATGCTAGACAGGGAGG
AGCTCTGTGGCTCCAATGAGCCTTGTGTCTGTATTTCCAAGTGTTAATGAAAAACCCACGCAGTTTTT
ACAAATTGAGCTCCAGGTCAGGGATATAAATGATCACTCTCCGCTTTCTTGGAAAAAGAAATGCTCTTA
GAAATCCAGAGAACAGTCCTGTTGGTGCTGTGTTCTTGCTTGAAGTGCAAAGGATTTAGATGAGGAA
TCAATGCTGTAAAAAGTACACAATAAATCCGAACCTCATTTCCAGTTAAAAAAGAGTCAATCCAGA
CAATAGGAAATACCTGAGTTAGTTCTGGACAAGGCGCTGGATTATGAAGAGCGCCCGAGCTCAGTTTC
ATCCTCACTGCTCTGGATGGCGGGTCCCCTCCAGGTCTGGAACCTGCCTGGTCAGGGTGGTGGTTGTAG
ATATTAATGACAACTCCCCTGAGTTTGAGCAGGCTTTTATGAGGTGAAGATTCTGGAGAATAGCATCCT
TGGTCCCCTGGTTGTGACCGTCTCAGCCTGGGATTTAGACTCTGGAACAAACAGTGAATATCCTATACC
TTTTCCCATGCCTCAGAAGATATTCGAAGACATTTGAAATTAATCAAAAGTCTGGTGACATTACTTTAA
CAGCACCTTTGGATTTTGAAGCAATTGAGTCATACTCAATAATCATTCAAGCCACAGATGGGGGAGGACT
TTTTGAAAAATCTACAGTCAGAATTCAGGTGATGGATGTAACGACAACGCTCCTGAAATCACTGTGTCA
TCAATTACAGTCCAATCCAGAAAACACTCCAGAGACTGTGGTTATGGTTTTTCAGGATACGAGACAGAG
ACTCTGGGGACAACGAAAGATGGTTTGTCTATCCCGAGGACATCCCATTCGTGCTAAAACTTCGGT
AAATAATTACTACACTTTGGAACAGAGAGACCGCTGGACAGAGAGAGCAGAGCCGAGTACAACATCACC
ATCACCGTCACCGACTTGGGGACCCCAAGGCTAAAAACCGAGCACAACATAACCGTGCTGGTCTCCGACG
TCAATGACAACGCCCGCCCTTACCCAACTTCTACGCCCTGTTCTGTCGCGAGAACAACAGCCCCGC
CCTGCACATCGGCAGCATCAGCGCCACAGACAGAGACTCGGGCACCACGCCAGGTCAACTACTCGTG
CTGCCGTCCAGGACCCGCACTGCCCTCGCCTCCCTGGTCTCCATCAACGCGGACAACGCCCACTGT
TTGCCCTCAGGTCGCTGGAATACGAGGCCCTGCAGGGTTCCAGTTCGCGTGGGCGCCACAGACCAGG
CTCCCCGGCTTTGAGCAGCGAGGCGCTGGTGCCTGCTGGTCTGGACGCCAACGACAACCTCGCCCTTC
GTGCTGTACCCGCTGCAGAACGGCTCCGCGCCCTGCACCGAGCTGGTGCCTGGGCGGCCGAGCCGGCT
ACCTGGTGACCAAGGTGGTGGCGGTGGACGGTGACTCGGGCCAGAACGCTGGCTGTGCTACCGAGTGT
CAAGGCCACGGAGCCCGGGCTATTGCGGTGTGGGCGCACAATGGCGAGGTGCGCACCGCCAGGCTGCTG
AGCGAGCGCGACGCGGCCAAGCACAGGCTGGTGGTGTGGTCAAGGACAATGGCGAGCTCCGCGCTCGG
CCACCGCCACGCTGCACGTGCTCCTGGTGGACGGCTTCTCCAGCCCTACCTGCCTCTCCCGAGGCGGC
CCCGGCCAGGCCAGGCCGACTCGCTCACTGTCTACCTGGTAGTGCGTTGGCCTCAGTGTGCTGCTGCTC
TTCTCTTCTCGGTGCTCCTGTTCTGGCGGTGCGGCTGTGCAGGAGGAGCAGGGCGGCCCGGTGCGTC
GCTGCTCGGTGCTGAGGGCCCTTTCCAGGACATCTGGTGGACGTGAGTGGACCGGGACCTGTCCCA
GAGCTACCACTATGAGGTGTGTGACTGGAGGCTCCAGGTCAAATAAGTTCAAATTTCTGAAACCAATT
ATCCCAACTTCTACCCAGAGCACAGGTAGTGAAGTGAAGAAAATCCCCATTTCAGAATAATTG
GTTTC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence:

>RG207172 representing NM_018932

Red=Cloning site Green=Tags(s)


MENGGAGTLQIRQVLLFFVLLGMSQAGSETGNFLVMEELQSGSFVGNLAKTLGLEVSELSSRGARVVSND
NKECLQLDTNTGDLRLREMLDREELCGSNEPCVLYFQVLMKNPTQFLQIELQVRDINDHSPVLEKEMLL
EIPENSPVGAVFLLSAKDLDVGINAVKSYTINPNSHFHVKIRVNPDRKYPELVLDKALDYEERPELSF
ILTALDGGSPPRSGTALVRVVVDINDNSPEFEQAFYEVKILENSILGSLVVTVSAWDLDSGTNSELST
FSHASEDIRKTFEINQKSGDITLTAPLDFAIESYSIIQATDGGGLFGKSTVRIQVMDVNDNAPEITVS
SITSPIPENTPETVVMVFRIRDRSDGNGKMCVIPEDIPFLKSSVNNYYTLETERPLDRESRAEYNIT
ITVTDLGTPLKTEHNITVLVSDVNDNAFTQTSYALFVRENNPALHIGSISATDRSDGTNAQVNSL
LPSQDPHLPLASLVSINADNGHLFALRSLDYELQGFQFRVGATDHGSPALSSEALVRVLVLDANDNSPF
VLYPLQNGSAPCTELVPWAAEPGYLVTKVVAVDGDSGQNAWLSYQLLKATEPGLFGVWAHNGEVRTARLL
SERDAAKHRLVVLVKDNGEPPRSATATLHVLLVDGFSQPYLPLPEAAPAQADSLTVYLVVALASVSSL
FLFVLLFVAVRLCRRRAAPVGRCSVPEGPFPHGLVDVSGTGTLQSQSYHYEVCVTGGSRSNKFKFLKPI
IPNPLPQSTGSEVEENPPFQNNLGF

TRTRPLE - GFP Tag - V

Restriction Sites:

SgfI-MluI

Cloning Scheme:



ACCN:

NM_018932

ORF Size:

2385 bp

OTI Disclaimer:


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_018932.3 , NP_061755.1
RefSeq Size:	3422 bp
RefSeq ORF:	2388 bp
Locus ID:	56124
UniProt ID:	Q9Y5F1
Cytogenetics:	5q31.3
Protein Families:	Transmembrane
Gene Summary:	This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RG207172