

Product datasheet for RG208687

ITM2C (NM 001012514) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: ITM2C (NM_001012514) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: ITM2C

Synonyms: BRI3; BRICD2C; E25; E25C; ITM3

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG208687 representing NM_001012514
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

GTGGTGGAGACGCTCATCTGCGGGGTGGTG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

ITM2C (NM_001012514) Human Tagged ORF Clone - RG208687

Protein Sequence: >RG208687 representing NM_001012514

Red=Cloning site Green=Tags(s)


MVKISFQPAVAGIKGDKADKASASAPAPASATEILLTPARLARDNFFRCGVLYEDSLSSQVRTQMELEED VKIYLDENYERINVPVPQFGGGDPADIIHDFQRGLTAYHDISLDKCYVIELNTTIVLPPRNFWELLMNVK RGTYLPQTYIIQEEMVVTEHVSDKEALGSFIYHLCNGKDTYRLRRRATRRRINKRGAKNCNAIRHFENTF VVETLICGVV


TRTRPLE - GFP Tag - V

Restriction Sites: Sg

Sgfl-Mlul

Cloning Scheme:





ACCN: NM_001012514

ORF Size: 660 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 001012514.3

RefSeq Size: 1982 bp
RefSeq ORF: 663 bp
Locus ID: 81618
UniProt ID: Q9NQX7
Cytogenetics: 2q37.1


Protein Families: Transmembrane

Gene Summary: Negative regulator of amyloid-beta peptide production. May inhibit the processing of APP by

blocking its access to alpha- and beta-secretase. Binding to the beta-secretase-cleaved APP C-terminal fragment is negligible, suggesting that ITM2C is a poor gamma-secretase cleavage inhibitor. May play a role in TNF-induced cell death and neuronal differentiation (By

similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RG208687