

Product datasheet for **RG208738**

RAN (NM_006325) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	RAN (NM_006325) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	RAN
Synonyms:	ARA24; Gsp1; TC4
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG208738 representing NM_006325 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCTGCGCAGGGAGAGCCCCAGGTCCAGTTCAACTTGTATTGGTTGGTGATGGTGGTACTGGAAAA
CGACCTTCGTGAAACGTCATTTGACTGGTGAATTTGAGAAGAAGTATGTAGCCACCTTGGGTGTTGAGGT
TCATCCCCTAGTGTCCACACCAACAGAGGACCTATTAAGTTCAATGTATGGACACAGCCGCCAGGAG
AAATTCGGTGGACTGAGAGATGGCTATTATATCCAAGCCAGTGTCCATCATAATGTTTGATGTAACAT
CGAGAGTTACTTACAAGAATGTGCCTAACTGGCATAGAGATCTGGTACGAGTGTGTGAAAACATCCCCAT
TGTGTTGTGTGGCAACAAGTGGATATTAAGGACAGGAAAGTGAAGGCGAAATCCATTGTCTCCACCGA
AAGAAGAATCTTCAGTACTACGACATTTCTGCCAAAAGTAACTACAACCTTGAAAAGCCCTTCTCTGGC
TTGCTAGGAAGCTCATTGGAGACCCTAAGTGGATTTGTTGCCATGCCTGCTCTCGCCCCACCAGAAGT
TGTGATGGACCCAGCTTTGGCAGCACAGTATGAGCACGACTTAGAGGTTGCTCAGACAACCTCTCCCC
GATGAGGATGATGACCTG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG208738 representing NM_006325
 Red=Cloning site Green=Tags(s)

MAAQGEPQVQFKLVLVGDGGTGKTTFVKRHLTGEFEKKYVATLGVEVHPLVFHTNRGPIKFNVWDTAGQE
 KFGGLRDGYIQAQCAIIMFDVTSRVTYKNVPNWHRDLVRVCENIPVLCGNKVDIKDRKVKAKSIVFHR
 KKNLQYYDISAKSNYNFEKPFLLWLRKLIIGDPNLEFVAMPALAPPEVVMDFALAAQYEHDLVAQTALP
 DEDDDL

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_006325

ORF Size: 648 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_006325.5
RefSeq Size:	1656 bp
RefSeq ORF:	651 bp
Locus ID:	5901
UniProt ID:	P62826
Cytogenetics:	12q24.33
Domains:	ras, RAN, RAS, RHO, RAB
Protein Families:	Druggable Genome, Transcription Factors
Gene Summary:	<p>RAN (ras-related nuclear protein) is a small GTP binding protein belonging to the RAS superfamily that is essential for the translocation of RNA and proteins through the nuclear pore complex. The RAN protein is also involved in control of DNA synthesis and cell cycle progression. Nuclear localization of RAN requires the presence of regulator of chromosome condensation 1 (RCC1). Mutations in RAN disrupt DNA synthesis. Because of its many functions, it is likely that RAN interacts with several other proteins. RAN regulates formation and organization of the microtubule network independently of its role in the nucleus-cytosol exchange of macromolecules. RAN could be a key signaling molecule regulating microtubule polymerization during mitosis. RCC1 generates a high local concentration of RAN-GTP around chromatin which, in turn, induces the local nucleation of microtubules. RAN is an androgen receptor (AR) coactivator that binds differentially with different lengths of polyglutamine within the androgen receptor. Polyglutamine repeat expansion in the AR is linked to Kennedy's disease (X-linked spinal and bulbar muscular atrophy). RAN coactivation of the AR diminishes with polyglutamine expansion within the AR, and this weak coactivation may lead to partial androgen insensitivity during the development of Kennedy's disease. [provided by RefSeq, Jul 2008]</p>

Product images:



Circular map for RG208738