

Product datasheet for RG208758

LIN7C (NM_018362) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Tag: TurboGFP

Symbol: LIN7C

Synonyms: LIN-7-C; LIN-7C; MALS-3; MALS3; VELI3

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide Sequence: >RG208758 representing NM_018362

Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGCCGCCTAGGGGAACCCGTGCGCTGGAGAGAGATATTTGTAGAGCAATTGAATTATTGGAAAAAC
TACAAAGGAGTGGAGAAGTACCACCACAGAAACTTCAGGCTTTGCAAAGAGTCCTTCAAAGTGAATTCTG
CAATGCTGTGAGAGAGGGTATATGAACATGTCTATGAGACTGTGGACATCAGTAGCAGTCCTGAAGTGAGA
GCGAACGCTACTGCAAAAGCCTACTGTTGCTGCATTTGCTGCCAGTGAAGGACATTCTCATCCTCGAGTTG
TTGAGCTACCAAAAACAGAAGAGGGCCTTGGATTCATATTTATGGGAGGCCAAAGAACAAACTCTCCAAT
CTATATATCCCGAATAATTCCAGGTGGAATTGCTGATAGACATGGGGGCCTCAAACGTGGAGATCAACTC
CTCTCTGTTAATGGAGTGAGTGTTGAAGGAGAACATCATGAAAAAGCTGTAGAACTGCTGAAAAGCCGCAC
AAGGAAAGGTTAAATTAGTGGTACGATACACCCCAAAGTCTTAGAAGAAAATGGAGTCGCGCTTTGAAAA

AATGAGATCAGCAAAACGCAGGCAACAGACC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG208758 representing NM_018362

Red=Cloning site Green=Tags(s)

MAALGEPVRLERDICRAIELLEKLQRSGEVPPQKLQALQRVLQSEFCNAVREVYEHVYETVDISSSPEVR ANATAKATVAAFAASEGHSHPRVVELPKTEEGLGFNIMGGKEQNSPIYISRIIPGGIADRHGGLKRGDQL

LSVNGVSVEGEHHEKAVELLKAAQGKVKLVVRYTPKVLEEMESRFEKMRSAKRRQQT

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn



Cloning Scheme:





ACCN: NM_018362

ORF Size: 591 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um

filter is required.

RefSeq: <u>NM_018362.4</u>

RefSeq Size: 4729 bp

RefSeq ORF: 594 bp

Locus ID: 55327

UniProt ID: Q9NUP9

Cytogenetics: llp14.1


Domains: PDZ, L27

Protein Families: Druggable Genome

Gene Summary: Plays a role in establishing and maintaining the asymmetric distribution of channels and

receptors at the plasma membrane of polarized cells. Forms membrane-associated multiprotein complexes that may regulate delivery and recycling of proteins to the correct membrane domains. The tripartite complex composed of LIN7 (LIN7A, LIN7B or LIN7C), CASK and APBAI may have the potential to couple synaptic vesicle exocytosis to cell adhesion in brain. Ensures the proper localization of GRIN2B (subunit 2B of the NMDA receptor) to neuronal postsynaptic density and may function in localizing synaptic vesicles at synapses where it is recruited by beta-catenin and cadherin. Required to localize Kir2 channels, GABA transporter (SLC6A12) and EGFR/ERBB1, ERBB2, ERBB3 and ERBB4 to the basolateral membrane of epithelial cells. [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RG208758