

## Product datasheet for **RG209344**

### CCL28 (NM\_148672) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** CCL28 (NM\_148672) Human Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** CCL28  
**Synonyms:** CCK1; MEC; SCYA28  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >RG209344 representing NM\_148672  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGCAGCAGAGAGGACTCGCCATCGTGGCCTTGCTGTCTGTGCGGCCCTACATGCCTCAGAAGCCATAC  
 TTCCATTGCCTCCAGCTGTTGCACGGAGGTTTCACATCATATTTCCAGAAGGCTCCTGAAAGAGTGAA  
 TATGTGTCGCATCCAGAGAGCTGATGGGATTGTGACTTGGCTGCTGTCATCCTTCATGTCAAGCGCAGA  
 AGAATCTGTGTCAGCCCGCACAACCACTACTGTTAAGCAGTGGATGAAAGTGCAAGCTGCCAAGAAAAATG  
 GTAAAGGAAATGTTTGCCACAGGAAGAAACACCATGGCAAGAGGAACAGTAACAGGGCACATCAGGGGAA  
 ACACGAAACATACGGCCATAAACTCCTTAT

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >RG209344 representing NM\_148672  
 Red=Cloning site Green=Tags(s)

MQQRGLAIVALVCAALHASEAILPIASSCCTEVSHHISRLLERVMCRIQRADGDCDLAAVILHVKRR  
 RICVSPHNHTVKQWMKVQAAKNGKGNVCHRKKHHGKRNSNRAHQKHETYGHKTPY

**TRTRPLE** - GFP Tag - V

**Restriction Sites:** Sgfl-MluI



[View online »](#)

**Cloning Scheme:**


**ACCN:** NM\_148672

**ORF Size:** 381 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.


**RefSeq:** [NM\\_148672.3](#), [NP\\_683513.1](#)

**RefSeq Size:** 816 bp

**RefSeq ORF:** 384 bp

**Locus ID:** 56477  
**UniProt ID:** [Q9NRJ3](#)  
**Cytogenetics:** 5p12  
**Protein Families:** Druggable Genome, Secreted Protein  
**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction  
**Gene Summary:** This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for resting CD4 or CD8 T cells and eosinophils. The product of this gene binds to chemokine receptors CCR3 and CCR10. This chemokine may play a role in the physiology of extracutaneous epithelial tissues, including diverse mucosal organs. Multiple transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Sep 2014]

### Product images:



Circular map for RG209344