



**Protein Sequence:** >RG209852 representing NM\_001001330  
 Red=Cloning site Green=Tags(s)

MVSWMISR AVL VFGMLYPAYYSYKAVKTKNVKEYVVRWMMYWIVFALYTVIETVADQTVAVWFPLYELKI  
 AFVIWLLSPYTKGASLIYRKFLHPLLSSKEREIDDIYVQAKERGYETMVNFGROGLNLAATAAVTAAVKS  
 QGAIATERLRSFSMHDLTITIQGDEPVGQRPYQPLPEAKKSKPAPSESAGYGIPLKDGDEKTDDEEAEGPYS  
 DNEMLTHKGLRRSQSMKSVKTTKGRKEVRYGSLKYKVKKRPQVYF

TRTRPLE - GFP Tag - V

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_001001330

**ORF Size:** 765 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** [NM\\_001001330.3](#)

**RefSeq Size:** 2308 bp

**RefSeq ORF:** 768 bp

**Locus ID:** 221035


**UniProt ID:** [Q6NUK4](#)

**Cytogenetics:** 10q21.3

**Protein Families:** Druggable Genome, Transmembrane

**Gene Summary:** Microtubule-binding protein required to ensure proper cell division and nuclear envelope reassembly by sequestering the endoplasmic reticulum away from chromosomes during mitosis. Probably acts by clearing the endoplasmic reticulum membrane from metaphase chromosomes.[UniProtKB/Swiss-Prot Function]

## Product images:



Circular map for RG209852