

Product datasheet for **RG210732**

HLA-DRB3 (NM_022555) Human Tagged ORF Clone

Product data:

| | |
|---------------------------|---|
| Product Type: | Expression Plasmids |
| Product Name: | HLA-DRB3 (NM_022555) Human Tagged ORF Clone |
| Tag: | TurboGFP |
| Symbol: | HLA-DRB3 |
| Synonyms: | DRB3; HLA-DPB1; HLA-DR1B; HLA-DR3B; HLA-DRB3* |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-AC-GFP (PS100010) |
| E. coli Selection: | Ampicillin (100 ug/mL) |
| ORF Nucleotide Sequence: | >RG210732 representing NM_022555 Red=Cloning site Blue=ORF Green=Tags(s) |

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGTGTGCTGAAGCTCCCTGGAGGCTCCAGCTTGGCAGCGTTGACAGTGACACTGATGGTGCTGAGCT
CCCGACTGGCTTTCGCTGGGGACACCCGACCACGTTTCTTGGAGCTGCTTAAGTCTGAGTGCATTTCTT
CAATGGGACGGAGCGGGTGGGTTCTTGGAGAGACTTCCATAACCAGGAGGAGTACGCGCGCTTCGAC
AGCGACGTGGGGAGTACCGGGCGGTGAGGGAGCTGGGGCGCCTGATGCCGAGTACTGGAACAGCCAGA
AGGACCTCCTGGAGCAGAAGCGGGCCAGGTGGACAATTACTGCAGACACAACACTACGGGGTTGGTGAGAG
CTTCACAGTGCAGCGCGAGTCCATCCTCAGGTGACTGTGTATCCTGCAAGACCCAGCCCTGCAGCAC
CACAACCTCCTGGTCTGCTCTGTGAGTGGTTTCTATCCAGGCAGCATTGAAGTCAGGTGGTTCGGAACG
GCCAGGAAGAGAAGGCTGGGGTGGTGTCCACGGCCTGATCCAGAATGGAGACTGGACCTTCAGACCCCT
GGTGTGCTAGAAACAGTTCTCGGAGTGGAGAGGTTTACACCTGCCAAGTGGAGCACCAAGCGTAACG
AGCCCTCTCACAGTGGAAATGGAGTGCACGGTCTGAATCTGCACAGAGCAAGATGCTGAGTGGAGTCGGGG
GCTTTGTGCTGGCCTGCTCTTCTTGGGGCCGGGCTGTTTCATCTACTTCAGGAATCAGAAAGGACACTC
TGGACTCAGCCAACAGGATTCTGAGC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG210732 representing NM_022555


Red=Cloning site Green=Tags(s)

MVCLKLPGGSSLAALTVTLMLVSSRLAFAGDTRPRFLELLKSECHFFNGTERTVRFLEHRHFHNQEEYARFD
SDVGEYRAVRELGRPDAEYWNQKDLLEQKRGQVDNYCRHNYGVGESFTVQRRVHPQVTVYPAKTQPLQH
HNLLVCSVSGFYPGSIEVRWFRNGQEEKAGVSTGLIQNGDWTFTLVMLETVPRSGEVYTCQVEHPSVT
SPLTVEWSARSESAQSKMLSGVGGFVLGLLFLGAGLFIYFRNQKGHSLQPTGFLS

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_022555

ORF Size: 798 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_022555.3](#), [NP_072049.2](#)

RefSeq Size: 1158 bp

RefSeq ORF: 801 bp

Locus ID: 3125

UniProt ID: [P79483](#)

Cytogenetics: 6p21.3


Domains: MHC_II_beta, ig, IGc1

Protein Families: Transmembrane

Protein Pathways: Allograft rejection, Antigen processing and presentation, Asthma, Autoimmune thyroid disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Hematopoietic cell lineage, Systemic lupus erythematosus, Type I diabetes mellitus, Viral myocarditis

Gene Summary: HLA-DRB3 belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells. The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. There are multiple pseudogenes of this gene. [provided by RefSeq, Feb 2020]

Product images:



Circular map for RG210732