

## **Product datasheet for RG211437**

# HNMT (NM 001024074) Human Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** HNMT (NM\_001024074) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: HNMT

Synonyms: HMT; HNMT-S1; HNMT-S2; MRT51

Mammalian Cell Neomycin

Selection:

**Vector:** pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG211437 representing NM\_001024074
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

CCAGAATTGCTGT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG211437 representing NM\_001024074

Red=Cloning site Green=Tags(s)

MASSMRSLFSDHGKYVESFRRFLNHSTEHQCMQEFMDKKLPGIIGRYQNCC

TRTRPLE - GFP Tag - V

**Restriction Sites:** Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




### **Cloning Scheme:**





#### Plasmid Map:



**ACCN:** NM\_001024074

ORF Size: 153 bp

#### HNMT (NM\_001024074) Human Tagged ORF Clone - RG211437

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 001024074.3</u>

 RefSeq Size:
 767 bp

 RefSeq ORF:
 156 bp

 Locus ID:
 3176

 UniProt ID:
 P50135

 Cytogenetics:
 2q22.1

Protein Families: Druggable Genome
Protein Pathways: Histidine metabolism

**Gene Summary:** In mammals, histamine is metabolized by two major pathways: N(tau)-methylation via

histamine N-methyltransferase and oxidative deamination via diamine oxidase. This gene encodes the first enzyme which is found in the cytosol and uses S-adenosyl-L-methionine as the methyl donor. In the mammalian brain, the neurotransmitter activity of histamine is controlled by N(tau)-methylation as diamine oxidase is not found in the central nervous system. A common genetic polymorphism affects the activity levels of this gene product in red blood cells. Multiple alternatively spliced transcript variants that encode different

proteins have been found for this gene. [provided by RefSeq, Jul 2008]