

## Product datasheet for **RG212080**

### **MYEOV2 (COPS9) (NM\_138336) Human Tagged ORF Clone**

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** MYEOV2 (COPS9) (NM\_138336) Human Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** COPS9  
**Synonyms:** CSNAP; MYEOV2  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >RG212080 representing NM\_138336  
**Red=Cloning site Blue=ORF Green=Tags(s)**

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGTGGCGCGCGCCGGAAGCGGCTCTGAGGCCGGAAGTGAGCCTAGAGCGCCGCGGCCCGAGATGAAGC  
CGGCGGTGGACGAGATGTTCCCCGAGGGCGCCGGGCCCTACGTGGACCTGGACGAGGTGGCACGGGCCCG  
GCGGGAGTCTCCGAGCGCGGGAGGCAGCACCGGGCTCTTGATGGACTTGGCAGCCAATGAAAAGGCCGT  
CATGCAGACTTTTTTAACGATTTTGAAGATCTTTTGTATGATGATGACATCCACAGCTCAGGACTGCCCA  
GGACAAGCCAGCAGTCTCCATGGTCCCGCACTGCAGCGAGGGCAGTCGGAGCACGGGGTCCGTGGGAA  
GGCAGCAGAGCGCCTGTGGTGTGCGGAAGAGAGATGTGAAGTGAACGGGAGAGAAGTGGCAGCTTTGGAC  
AGAGCGTTTGGGAGCACAGGCCATGGGCAGGGGCGGAGGCGCTAATGTTTACCAGATGCAGAGAATC  
CGCTGTGTGGCACCACAAGCAACAAGTGAAGGGGAAAATGGGCACAGGACGCTTAAGAAACAGTTTACG  
GAAGAACCAATCCAAGTGGCTGGGAAGTTATCTAGAGGTGCTCAGAACCACAAGAAGCAGGCGAGAGGTC  
AGCGAAGACAGCACAATATCAGTTTCTACTCACTGGAGGGGTAATGTTTAAAGAGTGATGAAACACCAT  
CCGTTGCTGGTGGAGAAGAAGGAAGAAGACCACACAGCCATGCATAGATGTGAGG

**ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA**



[View online »](#)

**Protein Sequence:** >RG212080 representing NM\_138336  
Red=Cloning site Green=Tags(s)

MWRAPEAALRPEVSLERRGPEMKPAVDEMFPPEGAGPYVDLDEVARARRESPSAGGSTGLLMDLAANEKAV  
HADFFNDFEDLFDDDDIHSSGLPRTSQSSMVPALQRGQSEHGVRGKAAERPVVSEERCELANGREVAALD  
RAFGSTGHGQGAELMFTRCREHPLCGTNKATSEGKMTGRLRNSLRKNQSKWLGSYLEVLRTRRSRREV  
SEDSTISVSTHWRGKCFKSDETPSVAGGEEGKKTTPCIDVR

TRTRPLE - GFP Tag - V


**Restriction Sites:** Sgfl-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shutting:



**Plasmid Map:**



**ACCN:** NM\_138336

**ORF Size:** 756 bp

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_138336.1</a> , <a href="#">NP_612209.1</a>
<b>RefSeq Size:</b>	759 bp
<b>RefSeq ORF:</b>	759 bp
<b>Locus ID:</b>	150678
<b>UniProt ID:</b>	<a href="#">Q8WXC6</a>
<b>Cytogenetics:</b>	2q37.3
<b>Gene Summary:</b>	Isoform 1: Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IκappaBα/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. Plays a role in cell proliferation. [UniProtKB/Swiss-Prot Function]