

## Product datasheet for **RG212720**

### ICAM2 (NM\_001099787) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	ICAM2 (NM_001099787) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	ICAM2
Synonyms:	CD102
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG212720 representing NM_001099787 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGTCCTCTTTTCGGTTACAGGACCCTGACTGTGGCCCTTCCACCCTGATCTGCTGTCCAGGATCGGATG  
AGAAGGTATTCGAGGTACACGTGAGGCCAAAGAAGCTGGCGGTTGAGCCCAAAGGGTCCCTCGAGGTCAA  
CTGCAGCACCACCTGTAACCAGCCTGAAGTGGTGGTCTGGAGACCTCTCTAGATAAGATTCTGCTGGAC  
GAACAGGCTCAGTGGAAACATTACTTGGTCTCAAACATCTCCCATGACACGGTCCCAATGCCACTTCA  
CCTGCTCCGGGAAGCAGGAGTCAATGAATTC AACGTGACGCGTGTACCAGCCTCCAAGGCAGGTCATCCT  
GACACTGCAACCCACTTTGGTGGCTGTGGCAAGTCCCTTACCATTGAGTGCAGGGTGCCACCGTGGAG  
CCCCTGGACAGCCTCACCTCTTCTGTTCCGTGGCAATGAGACTCTGCACTATGAGACCTTCGGGAAGG  
CAGCCCCCTGCTCCGACAGGAGGCCACAGCCACATTAACAGCACGGCTGACAGAGAGGATGGCCACCGCAA  
CTTCTCTGCCTGGCTGTGCTGGACTTGATGTCTCGCGGTGGCAACATCTTTCACAAACTCAGCCCCG  
AAGATGTTGGAGATCTATGAGCCTGTGTCGGACAGCCAGATGGTCATCATAGTCACGGTGGTGTCCGGTGT  
TGCTGTCCCTGTTGTCGATCTGTCTGCTCTGCTTTCATCTTCGGCCAGCACTTGGCCAGCAGCGGAT  
GGGCACCTACGGGTGCGAGCGGCTTGGAGGAGGCTGCCCCAGGCCTTCCGGCCA

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG212720 representing NM\_001099787  
 Red=Cloning site Green=Tags(s)

MSSFGYRTLTVLFTLICCPGSDEKVFVHVRPKKLAVEPKGSLEVNCSTTCNQPEVGGLETSLDKILLD  
 EQAQWKHYLVSNISHDTVLQCHF TCSGKQESMNSNVSVYQPPRQVILTLQPTLVAVGKSFTEICRVPTVE  
 PLDSLTLFLFRGNETLHYETFGKAAPAPQEATATFNSTADREDGHRNFSCLAVLDLMSRGGNIFHKHSAP  
 KMLEIYEPVSDSQMVIIIVTVSVLLSLFVTSVLLCFIFGQHLRQQRMGTYGVRAAWRRLLPQAFRP

TRTRPLE - GFP Tag - V


**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shutting:



**Plasmid Map:**



**ACCN:** NM\_001099787

**ORF Size:** 825 bp

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001099787.1</a> , <a href="#">NP_001093257.1</a>
<b>RefSeq Size:</b>	1270 bp
<b>RefSeq ORF:</b>	828 bp
<b>Locus ID:</b>	3384
<b>UniProt ID:</b>	<a href="#">P13598</a>
<b>Cytogenetics:</b>	17q23.3
<b>Protein Families:</b>	ES Cell Differentiation/IPS, Transmembrane
<b>Protein Pathways:</b>	Cell adhesion molecules (CAMs), Natural killer cell mediated cytotoxicity
<b>Gene Summary:</b>	The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008]