

## Product datasheet for **RG213823**

### **TAK1 (MAP3K7) (NM\_145332) Human Tagged ORF Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	TAK1 (MAP3K7) (NM_145332) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	MAP3K7
Synonyms:	CSCF; FMD2; MEKK7; TAK1; TGF1a
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>RG213823 representing NM\_145332  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGTCTACAGCCTCTGCCGCTCCTCCTCCTCGTCTTCGGCCGGTGAGATGATCGAAGCCCCTTCCC  
 AGGTCCTCAACTTTGAAGAGATCGACTACAAGGAGATCGAGGTGAAGAGTTGTTGGAAGAGGAGCCTT  
 TGGAGTTGTTTGCAAAGCTAAGTGGAGAGCAAAAGATGTTGCTATTAACAAATAGAAAAGTGAATCTGAG  
 AGGAAAGCGTTTATTGTAGAGCTTCGGCAGTTATCCCCTGTGAACCATCCTAATATTGTAAGCTTTATG  
 GAGCCTGCTTGAATCCAGTGTGCTTGTGATGGAATATGCTGAAGGGGGCTTTTATAATGTGCTGCA  
 TGGTGTGAACCATGCCATATTACTGCTGCCACGCAATGAGTTGGTGTTCACAGTGTCCCAAGGA  
 GTGGCTTATCTTACAGCATGCAACCCAAAGCGCTAATTCACAGGGACCTGAAACCACAACTTACTGC  
 TGGTTGCAGGGGGACAGTTCTAAAAATTTGTGATTTTGGTACAGCCTGTGACATTCAGACACACATGAC  
 CAATAACAAGGGGAGTGTGCTTGGATGCCACCTGAAGTTTTTGAAGGTAGTAATTACAGTGAAAAATGT  
 GAGCTCTTCAGCTGGGGTATTATTCTTTGGGAAGTGATAACGCGTCGAAACCCTTTGATGAGATTGGTG  
 GCCCAGCTTCCGAATCATGTGGGCTGTTCAATGGTACTCGACCACCTGATAAAAAATTTACCTAA  
 GCCCATTGAGAGCCTGATGACTCGTTGTTGGTCTAAAGATCCTTCCCAGCGCCCTTCAATGGAGGAAAT  
 GTGAAAAATAGACTCACTTGTGCGGTACTTTCCAGGAGCAGATGAGCCATTACAGTATCCTTGTGAGT  
 ATTCAGATGAAGGACAGAGCAACTCTGCCACCAGTACAGGCTCATTATGGACATTGCTTCTACAAATAC  
 GAGTAACAAAAGTGACACTAATATGGAGCAAGTTCTGCCACAAATGATACTATTAAGCGCTTAGAATCA  
 AAATTGTTGAAAAATCAGGCAAAGCAACAGAGTGAATCTGGACGTTTAAGCTTGGGAGCCTCCCGTGGGA  
 CGAGTGTGGAGAGCTTGCCCCCAACCTCTGAGGGCAAGAGGATGAGTGTGACATGCTGAAATAGAAGC  
 TAGGATCGCCGCAACCACAGCCTATTCCAAGCCTAAACGGGGCCACCGTAAAACTGCTTCATTTGGCAAC  
 ATTCTGGATGTCCCTGAGATCGTCATATCAGGCAACGGACAGCCAAGACGTAGATCCATCCAAGACTTGA  
 CTGTAACCTGGAACAGAACCTGGTCAGGTGAGCAGTAGGTCATCCAGTCCCAGTGTGAGAATGATTACTAC  
 CTCAGGACCAACCTCAGAAAAGCCAACCTCGAAGTCATCCATGGACCCCTGATGATTCCACAGATACCAAT  
 GGATCAGATAACTCCATCCCAATGGCTTATCTTACACTGGATCACCAACTACAGGCAAGAAGTGTGCA  
 GAACTGGACCAGGA

AC**GGGCCGCTCGAG** - GFP Tag - GTTTAA

**Protein Sequence:**

>RG213823 representing NM\_145332  
 Red=Cloning site Green=Tags(s)

MSTASAASSSSSSSAGEMIEAPSQVLNFFEEIDYKEIEVEEVVGRGAFGVVCKAKWRAKDVAIKQIESESE  
 RKAFIVELRQLSRVNHPIVVKLYGACLNVPCLVMEYAEGGSLYNVLHGAELPPYYTAAHAMSACLQCSQG  
 VAYLHSMQPKALIHRDLKPPNLLLVAAGTVLKIICDFGTACDIQTHMTNNKGSAAWMAPEVFEGSNYSEK  
 DVFSWGIILWEVITRRKPFDEIGGPAFRIMWAVHNGTRPPLIKNLPKPIESLMTRCWSKDPSQRPSMEEI  
 VKIMTHLMRYFPGADEPLQYPCQYSDEGQNSATSTGSFMDIASTNTSNKSDTNMEQVPATNDTIKRLS  
 KLLKNQAKQQSESGRLSLGASRGSSVESLPPTSEGKRMSADMSEIEARIAATTAYSKPKRGRHKATASFGN  
 ILDVPEIVISGNGQPRRRSIQDLTVTGTGTEPGQVSSRSSPSVRMITTSPTSEKPTRSHPWTPDDSDTN  
 GSDNSIPMAYLTLDHQLQARTSCRTGPG



TR**PLE** - GFP Tag - V

**Restriction Sites:**


Sgfl-NotI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



<b>ACCN:</b>	NM_145332
<b>ORF Size:</b>	1554 bp
<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_145332.3</a>
<b>RefSeq Size:</b>	2877 bp
<b>RefSeq ORF:</b>	1557 bp
<b>Locus ID:</b>	6885
<b>UniProt ID:</b>	<a href="#">O43318</a>
<b>Cytogenetics:</b>	6q15
<b>Protein Families:</b>	Druggable Genome, Protein Kinase
<b>Protein Pathways:</b>	Adherens junction, MAPK signaling pathway, NOD-like receptor signaling pathway, RIG-I-like receptor signaling pathway, T cell receptor signaling pathway, Toll-like receptor signaling pathway, Wnt signaling pathway
<b>Gene Summary:</b>	The protein encoded by this gene is a member of the serine/threonine protein kinase family. This kinase mediates the signaling transduction induced by TGF beta and morphogenetic protein (BMP), and controls a variety of cell functions including transcription regulation and apoptosis. In response to IL-1, this protein forms a kinase complex including TRAF6, MAP3K7P1/TAB1 and MAP3K7P2/TAB2; this complex is required for the activation of nuclear factor kappa B. This kinase can also activate MAPK8/JNK, MAP2K4/MKK4, and thus plays a role in the cell response to environmental stresses. Four alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]