

Product datasheet for **RG214035**

Glutathione Peroxidase 4 (GPX4) (NM_001039847) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Glutathione Peroxidase 4 (GPX4) (NM_001039847) Human Tagged ORF Clone
Symbol: Glutathione Peroxidase 4
Synonyms: GPx-4; GSHPx-4; MCSP; PHGPx; SMD5; snGPx; snPHGPx
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >RG214035 representing NM_001039847
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGAGCCTCGGCCGCTTTGCCGCTACTGAAGCCGGCGCTGCTCTGTGGGGCTCTGGCCGCGCCTGGCC
 TGGCCGGGACCATGTGCGCGTCCCGGGACGACTGGCGCTGTGCGCGCTCCATGCACGAGTTTTCCGCCAA
 GGACATCGACGGGCACATGGTTAACCTGGACAAGTACCGGGGCTTCGTGTGCATCGTCACCAACGTGGCC
 TCCAGTGAGGCAAGACCGAAGTAACTACACTCAGCTCGTGCACGCCGATACGCTGAGTGTG
 GTTTGCGGATCCTGGCCTTCCCGTGAACCACTCGGGAAGCAGGAGCCAGGGAGTAACGAAAGATCAA
 AGAGTTCGCCGCGGCTACAACGTCAAATTCGATATGTTAGCAAGATCTGCGTGAACGGGACGACGCC
 CACCCGCTGTGGAAGTGGATGAAGATCCAACCAAGGGCAAGGGCATCCTGGGAAATGCCATCAAGTGA
 ACTTCACCAAGTTTGGACACCGTCTCTCCACAGTTCCTCATCGACAAGAACGGCTGCGTGGTGAAGCGCT
 ACGGACCCATGGAGGAGCCCTGGTGTAGAGAAGGACCTGCCCACTATTTCTAGCTCCACAAGTGTG
 GGCCCCGCCGAGCCCTGCCACGCCCTTGGAGCCTTCCACCGCACTCA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG214035 representing NM_001039847
 Red=Cloning site Green=Tags(s)

MSLGRLCRLKLPALLCGALAAPLAGTMCASRDDWRCARSMHEFSAKDIDGHMVNLDKYRGFVCIVTNVA
 SQ*GKTEVNYTQLVDLHARYAECGLRILAFPCNQFGKQEPGSNEEIKEFAAGYNVKFDMFSKICVNGDDA
 HPLWKWMKIQPKGKILGNAIKWNFTKFGHRLSTVPHRQERLRGEALRTHGGAPGDREGPAPLFLAPQVC
 GPARAPAHALGAFHRHS

TRTRPLE - GFP Tag - V



[View online »](#)

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001039847

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#) The expression of this clone is not guaranteed due to the nature of selenoproteins.

OTI Annotation:

This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is encoded by UGA codon, which normally signals translational termination. Expression of this clone is not guaranteed due to the nature of selenoproteins.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001039847.3](#)

RefSeq Size: 964 bp

RefSeq ORF: 684 bp

Locus ID: 2879

UniProt ID: [P36969](#)


Cytogenetics: 19p13.3

Protein Families: Druggable Genome

Protein Pathways: Arachidonic acid metabolism, Glutathione metabolism

Gene Summary: The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyze the reduction of hydrogen peroxide, organic hydroperoxides and lipid hydroperoxides, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity. This isozyme has a high preference for lipid hydroperoxides and protects cells against membrane lipid peroxidation and cell death. It is also required for normal sperm development; thus, it has been identified as a 'moonlighting' protein because of its ability to serve dual functions as a peroxidase, as well as a structural protein in mature spermatozoa. Mutations in this gene are associated with Sedaghatian type of spondylometaphyseal dysplasia (SMDS). This isozyme is also a selenoprotein, containing the rare amino acid selenocysteine (Sec) at its active site. Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Transcript variants resulting from alternative splicing or use of alternate promoters have been described to encode isoforms with different subcellular localization. [provided by RefSeq, Dec 2018]

Product images:



Circular map for RG214035