

Product datasheet for **RG214679**

DIO2 (NM_013989) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
 Product Name: DIO2 (NM_013989) Human Tagged ORF Clone
 Symbol: DIO2
 Synonyms: 5DII; D2; DIOII; SELENOY; SelY; TXDI2
 Mammalian Cell Selection: Neomycin
 Vector: pCMV6-AC-GFP (PS100010)
 E. coli Selection: Ampicillin (100 ug/mL)
 ORF Nucleotide Sequence: >RG214679 representing NM_013989
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGGCATCCTCAGCGTAGACTTGCTGATCACACTGCAAATTCTGCCAGTTTTTTTCTCCAAGTGCCTCT
 TCCTGGCTCTCTATGACTCGGTCATTCTGCTCAAGCACGTGGTGTGCTGTTGAGCCGCTCCAAGTCCAC
 TCGCGGAGAGTGGCGGCGCATGCTGACCTCAGAGGGACTGCGCTGCGTCTGGAAGAGCTTCTCCTCGAT
 GCCTACAAACAGGTGAAATTGGGTGAGGATGCCCCCAATTCAGTGTGGTGCATGTCTCCAGTACAGAAG
 GAGGTGACAACAGTGGCAATGGTACCCAGGAGAAGATAGCTGAGGGAGCCACATGCCACCTCTTGACTT
 TGCCAGCCCTGAGCGCCCACTAGTGGTCAACTTTGGCTCAGCCACTTGACCTCCTTTCACGAGCCAGCTG
 CCAGCCTTCCGCAAACCTGGTGAAGAGTTCTCCTCAGTGGCTGACTTCTGCTGGTCTACATTGATGAGG
 CTCATCCATCAGATGGCTGGGCGATACCGGGGACTCCTCTTTGTCTTTGAGGTGAAGAAGCACCAGAA
 CCAGGAAGATCGATGTGCAGCAGCCAGCAGCTTCTGGAGCGTTTCTCCTTGCCGCCCCAGTGCCGAGTT
 GTGGCTGACCGCATGGACAATAACGCCAACATAGCTTACGGGGTAGCCTTTGAACGTGTGTGCATTGTGC
 AGAGACAGAAAATTGCTTATCTGGGAGGAAAGGGCCCTTCTCCTACAACCTTCAAGAAGTCCGGCATTG
 GCTGGAGAAGAATTCAGCAAGAGATGAAAGAAAAGTACTAGATTAGCTGGT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG214679 representing NM_013989
Red=Cloning site Green=Tags(s)

MGILSVDLLITLQILPVFFSNCLFLALYDSVILLKHVLLLLSRSKSTRGEWRRLTSEGLRCVWKSFLLD
 AYKQVKLGEDAPNSSVHVVSSTEGGDNNGTQEKIAEGATCHLLDFASPERPLVVNFGSAT*PPFTSQL
 PAFRKLVEEFSSVADFLLVYIDEAHPSDGWAIPGDSSLSFEVKKHQNQEDRCAAAQQLLERFSLPPQCRV
 VADRMNANIAYGVAFERVICVQRQKIAYLGGKGFPSYLNQEVRRHWLEKNFSKR*KKTRLAG

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_013989


OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#) The expression of this clone is not guaranteed due to the nature of selenoproteins.

OTI Annotation: This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is encoded by UGA codon, which normally signals translational termination. Expression of this clone is not guaranteed due to the nature of selenoproteins.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_013989.5</u>
RefSeq Size:	6703 bp
RefSeq ORF:	822 bp
Locus ID:	1734
UniProt ID:	<u>Q92813</u>
Cytogenetics:	14q31.1
Protein Families:	Druggable Genome
Gene Summary:	<p>The protein encoded by this gene belongs to the iodothyronine deiodinase family. It catalyzes the conversion of prohormone thyroxine (3,5,3',5'-tetraiodothyronine, T4) to the bioactive thyroid hormone (3,5,3'-triiodothyronine, T3) by outer ring 5'-deiodination. This gene is widely expressed, including in thyroid and brain. It is thought to be responsible for the 'local' production of T3, and thus important in influencing thyroid hormone action in these tissues. It has also been reported to be highly expressed in thyroids of patients with Graves disease, and in follicular adenomas. The intrathyroidal T4 to T3 conversion by this enzyme may contribute significantly to the relative increase in thyroidal T3 production in these patients. This protein is a selenoprotein containing the non-standard amino acid, selenocysteine (Sec), which is encoded by the UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Unlike the other two members (DIO1 and DIO3) of this enzyme family, the mRNA for this gene contains an additional in-frame UGA codon that has been reported (in human) to function either as a Sec or a stop codon, which can result in two isoforms with one or two Sec residues; however, only the upstream Sec (conserved with the single Sec residue found at the active site in DIO1 and DIO3) was shown to be essential for enzyme activity (PMID:10403186). Alternatively spliced transcript variants have been described for this gene. [provided by RefSeq, Oct 2018]</p>

Product images:



Circular map for RG214679