

Product datasheet for **RG215252**

DUSP4 (NM_057158) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	DUSP4 (NM_057158) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	DUSP4
Synonyms:	HVH2; MKP-2; MKP2; TYP
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG215252 representing NM_057158 Red=Cloning site Blue=ORF Green=Tags(s)

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGGAAGAAAAGTTCACTCCAACGGAAGCCAGTTTGCTGAACATAGCAGATCGCCAGGAGGACTGGGA
GAGACTGCAAACAGTTCGAGCCCCAGCATGGCGTTAGGTGTCAGCCAGCTGGCAGGAAGGTCCAGGTG
TCTGTGTTTCAGAGTCTCAAGGCGGCTATGAGAGGTTTTCTCCGAGTACCCAGAATTCTGTTCTAAAACC
AAGGCCCTGGCAGCCATCCACCCCGGTTCCCCCAGTGCCACAGAGCCCTTGACCTGGGCTGCAGCT
CCTGTGGGACCCACTACACGACCAGGGGGTCTGTGGAGATCCTTCCCTTCTCTACCTCGGCAGTGC
CTACCATGCTGCCCGGAGAGACATGCTGGACGCCCTGGGCATCACGGCTCTGTTGAATGTCTCTCGGAC
TGCCCAAACCACTTTGAAGGACACTATCAGTACAAGTGCATCCAGTGGAAGATAACCACAAGGCCGACA
TCAGCTCCTGGTTCATGGAAGCCATAGAGTACATCGATGCCGTGAAGGACTGCCGTGGGCGCGTGTGGT
GCACTGCCAGGCGGGCATCTCGCGGTCGGCCACCATCTGCCTGGCCTACCTGATGATGAAGAAACGGGTG
AGGCTGGAGGAGGCCCTTCGAGTTCGTTAAGCAGCGCCGAGCATCATCTCGCCAACTTCAGCTTCATGG
GGCAGCTGCTGCAGTTCGAGTCCCAGGTGCTGGCCACGTCCTGTGCTGCGGAGGCTGCTAGCCCCCTGGG
ACCCCTGCGGGAGCGGGCAAGACCCCGCCACCCCACTCGCAGTTTCGTCTTCAGCTTTCCGGTCTCC
GTGGGCGTGCACTCGGCCCCAGCAGCTGCCCTACCTGCACAGCCCCATCACCACTCTCCAGCTGT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)


Protein Sequence: >RG215252 representing NM_057158
 Red=Cloning site Green=Tags(s)

MGRKVVHNSQFAEHSRSPRRTGRDCKPVRAPSMALGVSQLAGRSRCLCSESQGGYERFSSEYPEFCST
 KALAAIPPPVPPSATEPLDLGCSSCGTPLHDQGGPVEILPFLYLGSAYHAARRDMLDALGITALLNVSSD
 CPNHFEHGYQYKCIPVEDNHKADISSWFMEAIEYIDAVKDCRGRVLVHCQAGISRSATICLAYLMMKKRV
 RLEEAFEFVKQRRSIIISPNFSFMGQLLQFESQVLATSCAAEAASPSGPLRERGKTPATPTSQFVFSFPVS
 VGVHSAPSSLPYLHSPITTSPTSC

TRTRPLE – GFP Tag – V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_057158

ORF Size: 909 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_057158.3</u>
RefSeq Size:	3404 bp
RefSeq ORF:	912 bp
Locus ID:	1846
UniProt ID:	<u>Q13115</u>
Cytogenetics:	8p12
Protein Families:	Phosphatase
Protein Pathways:	MAPK signaling pathway
Gene Summary:	<p>The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK1, ERK2 and JNK, is expressed in a variety of tissues, and is localized in the nucleus. Two alternatively spliced transcript variants, encoding distinct isoforms, have been observed for this gene. In addition, multiple polyadenylation sites have been reported. [provided by RefSeq, Jul 2008]</p>

Product images:



Circular map for RG215252