

Product datasheet for **RG217057**

P2Y6 (P2RY6) (NM_176796) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	P2Y6 (P2RY6) (NM_176796) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	P2RY6
Synonyms:	P2Y6
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG217057 representing NM_176796 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGAATGGGACAATGGCACAGGCCAGGCTCTGGGCTTGCCACCCACCACCTGTGTCTACCGGAGAACT
TCAAGCAACTGCTGCTGCCACCTGTGTATTCGGCGGTGCTGGCGGCTGGCCTGCCGCTGAACATCTGTGT
CATTACCCAGATCTGCACGTCCCGCCGGGCCCTGACCCGCACGGCCGTACACCCTAACCTTGCTCTG
GCTGACCTGCTATATGCCTGCTCCCTGCCCTGCTCATCTACAACATGCCCCAAGGTGATCACTGGCCCT
TTGGCGACTTCGCTGCCGCTGGTCCGCTTCTCTTCTATGCCAACCTGCACGGCAGCATCCTCTTCTCT
CACCTGCATCAGCTTCCAGCGCTACCTGGGCATCTGCCACCCGCTGGCCCCCTGGCACAACAGTGGGGGC
CGCCGGGCTGCCTGGCTAGTGTGTGTAGCCGTGTGGCTGGCCGTGACAACCCAGTGCCTGCCACAGCCA
TCTTCGCTGCCACAGGCATCCAGCGTAACCGCACTGTCTGCTATGACCTCAGCCCGCCTGCCCTGGCCAC
CCACTATATGCCCTATGGCATGGCTCTCACTGTATCGGCTTCTGCTGCCCTTTGCTGCCCTGTGGCC
TGCTACTGTCTCCTGGCCTGCCGCTGTGCCGCCAGGATGGCCCGGCAGAGCCTGTGGCCAGGAGCGGC
GTGGCAAGGGCGCCCGCATGGCCGTGGTGGTGGCTGCTGCCTTTGCCATCAGCTTCTGCCTTTTACAT
CACCAAGACAGCCTACCTGGCAGTGCCTCGACGCCGGCGTCCCTGCAGTGTATTGGAGGCCTTTGCA
GCGGCCTACAAAGGCACGCGCCGTTTCCAGTGCCAACAGCGTGCTGGACCCCATCCTCTTCTACTTCA
CCAGAAGAAGTTCGCCGGCGACACATGAGCTCCTACAGAACTCACAGCCAAATGGCAGAGGCAGGG
TCGC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG217057 representing NM_176796
Red=Cloning site Green=Tags(s)


MEWDNGTGQALGLPPTTCVYRENFKQLLLPPVYSAVLAAGLPLNICVITQICTSRRALTRTAVYTLNLAL
 ADLLYACSLPLLIYNYAQGDHWPFGDFACRLVRFLEYANLHGSILFLTCISFQRYLGICHPLAPWHKRGG
 RRAAWLVCVAVWLAVTTQCLPTAIFAATGIQRNRTVCYDLSPALATHYMPYGMALTVIGFLLPFAALLA
 CYLLACRLCRQDGAEPVAQERRGKAARMVAVVAAAF AISFLPFHITKTAYLAVRSTPGVPCTVLEAFA
 AAYKGRPFASANSVLDPIILFYFTQKKFRRRPHHELLQKL TAKWQRQGR

TRTRPLE - GFP Tag - V


Restriction Sites:

Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_176796

ORF Size: 984 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_176796.2
RefSeq Size:	1700 bp
RefSeq ORF:	987 bp
Locus ID:	5031
UniProt ID:	Q15077
Cytogenetics:	11q13.4
Protein Families:	Druggable Genome, GPCR, Transmembrane
Protein Pathways:	Neuroactive ligand-receptor interaction
Gene Summary:	The product of this gene belongs to the family of P2 receptors, which is activated by extracellular nucleotides and subdivided into P2X ligand-gated ion channels and P2Y G-protein coupled receptors. This family has several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor, which is a G-protein coupled receptor, is responsive to UDP, partially responsive to UTP and ADP, and not responsive to ATP. It is proposed that this receptor mediates inflammatory responses. Alternative splicing results in multiple transcript variants that encode different protein isoforms. [provided by RefSeq, Mar 2013]