

Product datasheet for **RG218310**

PSMA (FOLH1) (NM_004476) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PSMA (FOLH1) (NM_004476) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	PSMA
Synonyms:	FGCP; FOLH; GCP2; GCPII; mGCP; NAALAD1; NAALAdase; PSM; PSMA
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide
Sequence:

>RG218310 representing NM_004476
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGCATCGCC**

ATGTGGAATCTCCTTACGAAACCGACTCGGCTGTGGCCACCGCGCCGCCCGCGCTGGCTGTGCGCTG
GGGCGCTGGTGTGGCGGTGGCTTCTTCTCCTCGGCTTCTCTTCGGGTGTTTATAAAATCCTCCAA
TGAAGCTACTAACATTACTCCAAAGCATAATATGAAAGCATTCTTGGATGAATTGAAAGCTGAGAACATC
AAGAAGTCTTATATAATTTTACACAGATACCACATTTAGCAGGAACAGAACAAAATTTTTCAGCTTGCAA
AGCAAATTCATCCAGTGGAAGAATTTGGCCTGGATTCTGTTGAGCTAGCACATTATGATGTCCTGTT
GTCCTACCCAAATAAGACTCATCCCACTACATCTCAATAATTAATGAAGATGGAATGAGATTTTCAAC
ACATCATTATTTGAACCACCTCTCCAGGATATGAAAATGTTTCGGATATTGTACCACCTTTCAGTGCTT
TCTCTCCTCAAGGAATGCCAGAGGGCGATCTAGTGTATGTTAACTATGCACGAACTGAAGACTTCTTTAA
ATTGGAACGGGACATGAAAATCAATTGCTCTGGGAAAATTGTAATTGCCAGATATGGGAAAGTTTTCAGA
GAAAATAAGGTTAAAAATGCCAGCTGGCAGGGGCCAAAAGGAGTCACTCTACTCCGACCTGCTGACT
ACTTTGCTCCTGGGGTGAAGTCCTATCCAGATGTTTGAATCTTCTGGAGGTGGTGTCCAGCGTGGAAA
TATCCTAAATCTGAATGGTGCAGGAGACCTCTCACACCAGGTTACCCAGCAAATGAATATGCTTATAGG
CGTGGAATTGCAGAGGCTGTTGGTCTTCCAAGTATTCCTGTTTATCCAATTGGATACTATGATGCACAGA
AGCTCCTAGAAAAATGGGTGGCTCAGCACCACAGATAGCAGCTGGAGAGGAAGTCTCAAAGTGCCTTA
CAATGTTGGACCTGGCTTACTGGAACTTTTCTACACAAAAGTCAAGATGCACATCCACTCTACCAAT
GAAGTGACAAGAATTTACAATGTGATAGTACTCTCAGAGGAGCAGTGGAAACAGACAGATATGTCATTC
TGGGAGGTCACCGGGACTCATGGGTGTTTGGTGGTATTGACCCTCAGAGTGGAGCAGCTGTTGTTTATGA
AATTGTGAGGAGCTTTGGAACACTGAAAAGGAAGGGTGGAGACCTAGAAGAACAATTTTGTTCGAAGC
TGGGATGCAGAAGAATTTGGTCTTCTGGTCTACTGAGTGGGCAGAGGAGAATTCAGACTCCTTCAAG
AGCGTGGCGTGGCTTATATTAATGCTGACTCATCTATAGAAGGAACTACACTCTGAGAGTTGATTGTAC
ACCGCTGATGTACAGCTTGGTACACAACCTAACAAAAGAGCTGAAAAGCCCTGATGAAGGCTTTGAAGGC
AAATCTCTTTATGAAAGTTGGACTAAAAAAGTCTTCCCAAGAGTTCAGTGGCATGCCAGGATAAGCA
AATTGGGATCTGGAAATGATTTTGGAGTGTCTTCCAACGACTTGAATGCTTCAGGCAGAGCACGGTA
TACTAAAAATTTGGAAACAAACAATTCAGCGGCTATCCACTGTATCACAGTGTCTATGAAACATATGAG
TTGGTGGAAAAGTTTTATGATCCAATGTTTAAATATCACCTCACTGTGGCCAGGTCGAGGAGGGATGG
TGTTTGAGCTAGCCAATCCATAGTGTCCCTTTTGGATTGTCGAGATTATGCTGTAGTTTTAAGAAAGTA
TGCTGACAAAATCTACAGTATTTCTATGAAACATCCACAGGAAATGAAGACATACAGTGTATCATTGAT
TCACTTTTTCTGCAGTAAAGAATTTTACAGAAATGCTTCCAAGTTCAGTGAGAGACTCCAGGACTTTG
ACAAAAGCAACCAATAGTATTAAGAATGATGAATGATCAACTCATGTTTCTGGAAAGAGCATTTATTGA
TCCATTAGGGTTACCAGACAGGCCTTTTATAGGCATGTCATCTATGCTCCAAGCAGCCACAACAAGTAT
GCAGGGGAGTCATTCCCAGGAATTTATGATGCTCTGTTTATGATTGAAAGCAAAGTGGACCCTTCCAAGG
CCTGGGAGAAAGTGAAGAGACAGATTTATGTTGCAGCCTTACAGTGCAGGCAGCTGCAGAGACTTTGAG
TGAAGTAGCC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG218310 representing NM_004476
 Red=Cloning site Green=Tags(s)

MWLLHETDSAVATARRPRWLCAGALVLAGGFFLLGFLFGWFIKSSNEATNITPKHNMKAFLDELKAENI
 KKFLYNFTQIPHLAGTEQNFQLAKQIQSQWKEFGLDSVELAHYDVLLSYPNKTHPNYISIIINEDGNEIFN
 TSLFEPPIPGYENVSDIVPPFSAFSPQGMPEGLVYVNYARTEDDFKLERDMKINCSGKIVIARYGKVF
 GNKVNAQLAGAKGVILYSDPADYFAPGVKSYPDGWNLPGGGVQRGNILNLNGAGDPLTPGYPANEYAYR
 RGIAEAVGLPSIPVHPIGYDDAQKLEKMGGSAPPDSSWRGSLKVPYNVGPFTGNFSTQKVKMHIHSTN
 EVTRIYNVIGTLRGAVEPDRYVILGGHRDSWVFGGIDPQSGAAVVHEIVRSFGTLKKEGWRPRRILFAS
 WDAEEFLLGSTEWAEEENSRLQLQERGVAYINADSSIEGNYTLRVDCTPLMYSLVHNLTKELKSPDEGFEG
 KSLYESWTKKSPSEFSGMPRIKLGSGNDFEVFFQRLGIASGRARYTKNWETNKFSGYPLYHSVYETYE
 LVEKFYDPMFKYHLTVAQVRGGMVFEANISIVLPFDCRDYAVVLRKYADKIYSISMKHPQEMKTVSVSFD
 SLFSAVKNFTEIASKFSERLQDFDKSNPVLMMNDQLMFLERAFIDPLGLPDRPFYRHVIYAPSSHNKY
 AGESFPGIYDALFDIESKVDPSKAWGEVKRQIYVAaftvQAAAETLSEVA

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_004476

ORF Size: 2250 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_004476.3
RefSeq Size:	2653 bp
RefSeq ORF:	2253 bp
Locus ID:	2346
UniProt ID:	Q04609
Cytogenetics:	11p11.12
Domains:	PA, TFR_dimer, Peptidase_M28
Protein Families:	Druggable Genome, Protease, Transmembrane
Gene Summary:	<p>This gene encodes a type II transmembrane glycoprotein belonging to the M28 peptidase family. The protein acts as a glutamate carboxypeptidase on different alternative substrates, including the nutrient folate and the neuropeptide N-acetyl-l-aspartyl-l-glutamate and is expressed in a number of tissues such as prostate, central and peripheral nervous system and kidney. A mutation in this gene may be associated with impaired intestinal absorption of dietary folates, resulting in low blood folate levels and consequent hyperhomocysteinemia. Expression of this protein in the brain may be involved in a number of pathological conditions associated with glutamate excitotoxicity. In the prostate the protein is up-regulated in cancerous cells and is used as an effective diagnostic and prognostic indicator of prostate cancer. This gene likely arose from a duplication event of a nearby chromosomal region. Alternative splicing gives rise to multiple transcript variants encoding several different isoforms. [provided by RefSeq, Jul 2010]</p>

Product images:



Circular map for RG218310